

A GEOMETRICAL TREATISE
ON
CONIC SECTIONS.



A
GEOMETRICAL TREATISE
ON
CONIC SECTIONS.

WITH A COPIOUS COLLECTION OF EXAMPLES,
EMBODYING EVERY QUESTION WHICH HAS BEEN PROPOSED IN
THE SENATE HOUSE AT CAMBRIDGE.

For the Use of Schools and Students in the Universities.

BY THE REV. W. H. DREW, M.A.

ST. JOHN'S COLLEGE, CAMBRIDGE,

SECOND MASTER OF BLACKHEATH PROPRIETARY SCHOOL.

Second Edition.

Cambridge :
MACMILLAN AND CO.

MDCCLXII.

LONDON
PRINTED BY R. CLAY, SON, AND TAYLOR, BREAD STREET HILL

PREFACE

TO THE SECOND EDITION.

THE limits which I have assigned to myself both in this and the former edition, and which I have endeavoured not to transgress either in excess or defect, are the requirements of the Senate-House Examination at Cambridge. After a very careful revisal, I have come to the conclusion that the Propositions which I have given are fully sufficient for the solution of all such questions as can be conveniently treated geometrically. At the same time there is no proposition which can be omitted without the risk of some important point being overlooked. Though the alterations in the present edition are not extensive, they are important; one or two of the demonstrations have been much shortened and simplified, a footnote has been inserted on the first proposition both of the Ellipse and of the Hyperbola, where the text seemed obscure; a fresh proposition has been introduced on the circle of curvature in the Parabola and Ellipse, which is also applicable to the

Hyperbola; the letters of some of the figures have been changed to avoid the confusion which arose from the multiplication of dashes and suffixes; many corollaries have been added with a view of calling attention to points which had been passed over without notice. Finally, 130 fresh problems, selected from Examination Papers of recent dates, have been appended, which it is believed will afford the student all the opportunities for practising himself in this subject which he can desire.

The solutions of the problems contained in the former edition have been published in such a form as to be of service not only to the teacher, but also to the pupil. The additional problems, which are not referred to in the solutions, are placed separately at the end of the book. With the solutions the book now consists of three parts—the book-work, problems solved by way of illustration, and unsolved problems for the independent exercise of the student.

W. IL. DREW.

BLACKHEATH PROPRIETARY SCHOOL,

February 18th, 1862.

P R E F A C E.

THE importance of studying the properties of the Conic Sections by Geometrical methods before entering upon the algebraic theory cannot, I think, be too much insisted upon, whether regarded as an introduction to the study of Newton, or with a view to a more complete understanding subsequently of the principles of Analytical Geometry.

The advantage of observing this order is now so fully recognised at Cambridge, that during the first three days of the Examination for Honours the geometrical method is alone admissible.

At the same time I believe that this mode of treating the subject has not received in Schools the attention which it deserves, and I cannot but think that this is in a great degree owing to the want of a book altogether suited for the purpose.

In the following pages, which were originally compiled for the use of my own pupils at Blackheath,

I have aimed at supplying this deficiency, and I have endeavoured to place the subject before the student in such a form, that, after mastering the elements of Euclid, he may find it an easy and interesting continuation of his geometrical studies.

With a view also of rendering the work a complete Manual of what is required at Cambridge, I have either embodied into the text, or inserted among the examples, every book-work question, problem, and rider, which has been proposed in the Senate House up to the present time.

The principal points in which the present Treatise will be found to differ from those now in use are the following :—

(1.) The three Conic Sections are defined in a uniform manner.

(2.) The use of the Second Book of Euclid is avoided, *as much as possible*, as having a tendency to lead to algebraical methods of reasoning.

(3.) The properties of the ellipse and hyperbola which depend upon the directrix are fully given, and the analogy between these curves and the parabola is thus maintained.

(4.) A method of proving the fundamental property of the tangents has been adopted, which, while it presents the idea of a limit in its simplest form, is

applicable, word for word, to each* of the three curves.

(5.) With a view to simplification, several of the demonstrations have been much modified from the form in which they are usually presented, among which I may particularly mention those propositions which relate to the properties of conjugate diameters in the ellipse, and of the asymptotes in the hyperbola, and the proof that $QV^2 = 4SP \cdot PV$ in the parabola; while many important theorems admitting of an easy geometrical solution are introduced.

(6.) A full discussion is given in a distinct* chapter of the Sections of the Cone. Figures are drawn representing the position of the foci and of the directrices of the sections in every case.

W. H. DREW.

BLACKHEATH PROPRIETARY SCHOOL,

March 19th, 1857.


CONIC SECTIONS.

INTRODUCTION.

1. DEF. The curve traced out by a point, which moves in such a manner that its distance from a given fixed point continually bears the same ratio to its distance from a given fixed line, is called a *Conic Section*.

The fixed point is called the *Focus*, and the fixed line the *Directrix*.

Thus if S be the focus, and KK' the directrix, and P a point from which PM is drawn at right angles to the directrix, the curve traced out by P will be a *Conic Section*, provided P move in such manner that SP always bears the same ratio to PM .



(1.) When the distance from the fixed point is equal to the distance from the fixed line, that is, when SP is equal to PM , the *Conic Section* is called a *Parabola*.

(2.) When the distance from the fixed point is less than the distance from the fixed line, that is, when the ratio which

SP bears to PM is less than unity, the *Conic Section* is called an *Ellipse*.

(3.) When the distance from the fixed point is greater than the distance from the fixed line, that is, when the ratio which SP bears to PM is greater than unity, the *Conic Section* is called an *Hyperbola*.

2. The reason of the term *Conic Sections* being applied to these curves is that, when a *Cone* is intersected by a plane surface, the boundary of the section so formed will, *in general*, be one or other of these curves.

I purpose to investigate the properties of the *Conic Sections* from the definitions given above, and afterwards to show in what manner a *Cone* must be divided by a plane in order that the curve of intersection may be a *Parabola*, *Ellipse*, or *Hyperbola*.




CHAPTER I.

THE PARABOLA.

PROP. I.

3. THE focus and directrix of a parabola being given, to find any number of points on the curve.



Let S be the focus, and KK' the directrix.

Draw XSx at right angles to the directrix, and bisect the line SX in A ; then

$$\text{since } AS = AX,$$

$\therefore A$ is a point on the curve.

The point A is called the *Vertex*, and the line Ax , with respect to which the curve is evidently symmetrical, is called the *Axis*.

On the directrix take any point M ; join SM ; and draw MP at right angles to the directrix.

At the focus S make the angle MSP equal to the angle SMP ; then


$$SP = PM,$$

$\therefore P$ is a point on the curve.

So by taking any number of points, M' , M'' , on the directrix, we may obtain as many points, P' , P'' , on the curve as we please, and the line which passes through A and all these points will be the parabola whose focus is S and directrix KK' .

COR. 1. As M is taken further away from the point X , the line SM and the angles SMP , MSP , and, consequently, the lines SP and PM , continually increase. Hence, since XM and MP increase together, the curve recedes at the same time both from the axis and directrix; and since the angle SMP can never exceed a right angle, and the lines SP and MP will therefore always meet, it is evident that there is no limit to the distance to which the curve may extend on both sides of the axis.

COR. 2. The parabola may be described practically in the following manner.



Let S be the focus and KX be the directrix; and let a rigid bar QM , having a string of the same length as itself fastened at one end Q , be made to slide parallel to the axis with the other end M on the directrix; then if the other end of the string be fastened at the focus, and the string be kept stretched by means of the point of a pencil at P , in contact with the bar, since SP will always be equal to PM , it is evident that the point P will trace out the parabola.

PROP. II.

4. The distance of any point inside the parabola from the focus is less than its distance from the directrix; and the distance of any point outside the parabola from the focus is greater than its distance from the directrix.



(1.) Let Q be a point inside the parabola.

Draw QM at right angles to the directrix, meeting the parabola in P ; join SP ; then

$$\text{since } SP = PM,$$

$$\therefore SP \text{ and } PQ = QM.$$

$$\text{But } SP \text{ and } PQ > SQ,$$

$$\therefore QM > SQ.$$

(2.) Let Q be a point outside the parabola.

Draw MQ at right angles to the directrix, and produce it to meet the parabola in P ; join SP ; then

$$\therefore \text{ since } SQ \text{ and } QP > SP,$$

$$\text{and } SP = PM,$$

$$\therefore SQ \text{ and } QP > PM,$$


$$\therefore SQ > QM.$$

COR. Conversely a point will be inside or outside the parabola according as its distance from the focus is less or greater than its distance from the directrix.

5. DEF. The line PN (see fig. Prop. III.) drawn at right angles to the axis from the point P in the curve is called the *Ordinate* of the point P , and the line AN the *Abscissa*. The double ordinate BC drawn through the focus, and terminated both ways by the curve, is called the *Latus Rectum*.

PROP. III.

The Latus Rectum $BC = 4AS$.



Draw BK at right angles to the directrix.

Then $SB = BK = SX = 2AS$,


$\therefore BC = 4AS$.

6. DEF. If a point P' be taken on the ~~parabola~~^{curve} (see fig. *Prop. IV.*) near to P , and PP' be joined, the line PP' produced, in the limiting position which it assumes when P' is made to approach indefinitely near to P , is called the *Tangent* to the parabola at the point P .

PROP. IV.

• If the tangent to the parabola at any point P intersect the directrix in the point Z ; then SZ will be at right angles to SP .

Let P' be a point on the parabola near to P .



Draw the chord PP' , and produce it to meet the directrix in Z ; join SZ .

Draw PM , $P'M'$ at right angles to the directrix; join SP , SP' ; and produce PS to meet the parabola in Q .

Then, since the triangles ZMP , $ZM'P'$ are similar,

$$\begin{aligned}\therefore ZP : ZP' &:: MP : M'P', \\ &:: SP : SP',\end{aligned}$$

$\therefore SZ$ bisects the angle $P'SQ$. (*Euclid*, VI. *Prop.* A.)

Now when P' is indefinitely near to P , and PP' becomes the tangent at the point P , the angle $PS P'$ becomes indefinitely small, while the angle QSP' approaches two right angles, and therefore the angle $P'SZ$, which is half of the angle $P'SQ$, becomes ultimately a right angle.


Hence, when PZ is the tangent,

the angle ZSP is a right angle,
or SZ is perpendicular to SP .

COR. Conversely, if SZ be drawn at right angles to SP , meeting the directrix in Z , and PZ be joined, PZ will be a tangent at P .

PROP. V.

7. The tangent at any point P of a parabola bisects the angle between the focal distance SP , and the perpendicular PM on the directrix.



Let the tangent at P meet the directrix in the point Z ; join SZ ; then since the angle ZSP is a right angle, (*Prop. IV.*)

$$\therefore ZS^2 + SP^2 = PZ^2.$$

$$\text{Also } ZM^2 + MP^2 = PZ^2,$$

$$\therefore ZS^2 + SP^2 = ZM^2 + MP^2.$$

$$\text{But } SP = PM,$$

$$\therefore ZS = ZM.$$

Now in the triangles ZPS , ZPM ,

$$\therefore ZP, PS = ZP, PM, \text{ each to each.}$$

$$\text{and } ZS = ZM,$$

$$\therefore \text{the angle } SPZ = \text{the angle } MPZ;$$

$$\text{or } PZ \text{ bisects the angle } SPM.$$

COR. 1. If ZP be produced to R , then the angle $SPR =$ the angle MPR .

COR. 2. It is evident that the tangent at the vertex A is perpendicular to the axis.

PROP. VI.

8. The tangents at the extremities of a focal chord intersect at right angles in the directrix.

Let PSQ be a focal chord, and let the tangent at P meet the directrix in Z .

Join SZ ; then

the angle ZSP is a right angle, (*Prop. IV.*)

and $\therefore$ also the angle ZSQ is a right angle,

$\therefore ZQ$ is the tangent at Q , (*Prop. IV. Cor.*)

or the tangents at the extremities of the focal chord PSQ intersect in the directrix.

Again, draw PM, QM' at right angles to the directrix; then

since $MP, PZ = SP, PZ$, each to each,

and the angle $MPZ =$ the angle SPZ ,

$\therefore$ the angle $MZP =$ the angle SZP ,

$\therefore$ the angle SZP is half of the angle SZM .

So the angle SZQ is half of the angle SZM' ,

$\therefore$ the angle PZQ is half of the two SZM and SZM' .


But the angles SZM and $SZM' =$ two right angles,

$\therefore$ the angle PZQ is a right angle,

or the tangents at the extremities of a focal chord intersect *at right angles* in the directrix.

PROP. VII.

9. If the tangent at any point P of a parabola meet the axis produced in the point T , and PN be the ordinate of the point P , then $NT = 2AN$.



Join SP , and draw PM at right angles to the directrix; then

$\therefore$ the angle $SPT =$ the angle $MPT =$ the angle STP ,

$$\therefore ST = SP.$$

$$\text{But } SP = PM = XN,$$

$$\therefore ST = XN.$$

$$\text{But } AS = AX,$$

$$\therefore \text{the remainder } AT = \text{the remainder } AN,$$

$$\therefore NT = 2AN.$$

DEF. The line NT is called the *Subtangent*.

10. DEF. The line PG , drawn at right angles to PT , is called the *Normal* at the point P , and NG the *Subnormal*.

PROP. VIII.

If the normal at the point P of a parabola meet the axis in the point G , then $NG = 2AS$.

Since the angle SPG = the complement of the angle SPT ,
 and the angle $S\cancel{PG}$ = the complement of the angle STP ,
 and also the angle SPT = the angle STP , (*Prop. VII.*)

$\therefore$ the angle SPG = the angle SGP ,

$\therefore SG = SP$.

But $SP = PM = XN$,

$\therefore SG = XN$.

Taking away the common part SN ,

the remainder $NG = SX = 2AS$.

PROP. IX.

11. If PN be an ordinate to the parabola at the point P ;
 then $PN^2 = 4AS \cdot AN$.

Since TPG is a right angle, and PN perpendicular to TG ,

$\therefore PN$ is a mean proportional between TN and NG ;

or $PN^2 = TN \cdot NG$. (*Euclid, VI. 8 Cor.*)

But $TN = 2AN$, (*Prop. VII.*)

and $NG = 2AS$. (*Prop. VIII.*)

$\therefore PN^2 = 4AS \cdot AN$.

PROP. X.

12. If the tangent at any point P intersect the tangent at the vertex in Y , SY will bisect PT at right angles, and will be a mean proportional between SA and SP .

Draw PN at right angles to the axis; then

since AY is parallel to PN ,

$\therefore TY : YP :: TA : AN$.

But $AT = AN$, (*Prop. VII.*)

$\therefore TY = PY$;

and $\therefore SY, YP = SY, YT$, each to each,

and $SP = ST$, (*Prop. VII.*)

$\therefore$ the angle $SY P$ = the angle $SY T$,

$\therefore SY$ is perpendicular to PT .

CONIC SECTIONS.

Let O be the given external point.

Join OS , and with centre O and radius OS describe a circle, cutting the directrix in M and M' , which it will always do, on whichever side of the directrix O is situated, since O is nearer to the directrix than to the focus. (*Prop. II.*)

Draw MQ and $M'Q'$ parallel to the axis meeting the parabola in Q and Q' .

Join OQ , OQ' ; these will be the tangents required.

Join SQ and SQ' ; then

$$\begin{aligned} \therefore OQ, QS &= OQ, QM, \text{ each to each,} \\ \text{and } OS &= OM, \end{aligned}$$

$$\therefore \text{ the angle } OQS = \text{ the angle } OQM,$$

$$\therefore OQ \text{ is the tangent at } Q. \quad (\textit{Prop. V.})$$

$$\text{So } OQ' \text{ is the tangent at } Q'.$$

PROP. XII.

14. If from a point O a pair of tangents OQ and OQ' be drawn to a parabola, the triangles OSQ , OSQ' will be similar, and OS will be a mean proportional between SQ and SQ' .

Join SM , cutting OQ at right angles (*Prop. X. Cor.*) in the point Y ; then

$$\text{since the angle } SQO = \text{ the angle } MQO, \quad (\textit{Prop. V.})$$

$$\text{and that the angle } MQO = \text{ the angle } SMM',$$

each of these angles being the complement of the angle QMY ,

$$\therefore \text{ the angle } SQO = \text{ the angle } SMM'.$$

But the angle SMM' at the circumference is half the angle SOM at the centre, and is therefore equal to the angle SOQ .

$$\therefore \text{ the angle } SQO = \text{ the angle } SOQ.$$

$$\text{So the angle } SOQ = \text{ the angle } SQ'O,$$

$$\therefore \text{ the remaining angle } OSQ = \text{ the remaining angle } OSQ'.$$

And therefore the triangle OSQ is similar to the triangle OSQ' ,

$$\therefore SQ : SO :: SO : SQ',$$


$$\therefore SQ \cdot SQ' = SO^2,$$

or SO is a mean proportional between SQ and SQ' .

$$OQ^2 : OQ'^2 :: SQ : SQ', \text{ \& \& } \angle SOQ : \angle SOQ' :: SQ : SQ',$$

PROP. XIII.

15. If a pair of tangents OQ , OQ' be drawn to a parabola, and OV be drawn parallel to the axis meeting QQ' in V , then QQ' shall be bisected in V .



Draw QM , $Q'M'$ at right angles to the directrix.

Join OM , OM' ; and let OV meet MM' in Z .

Then, since $OM = OM'$, (*Prop. XI.*)

$\therefore$ the angle $OMZ =$ the angle $OM'Z$,

and the angle $OZM =$ the angle OZM' ,

and the side OZ is common to the triangles OZM , OZM' ,

$$\therefore MZ = M'Z.$$

And because the lines QM , ZV , $Q'M'$ are parallel,

$$\therefore QV : Q'V :: MZ : M'Z.$$


$$\text{But } MZ = M'Z,$$

$$\therefore QV = Q'V,$$

$$\therefore QQ' \text{ is bisected in } V.$$

PROP. XIV.

16. If from a point O a pair of tangents OQ , OQ' be drawn to a parabola, and OV be drawn parallel to the axis meeting the parabola in P , and QQ' in V , then the tangent at P will be parallel to QQ' , and OV will be bisected in P .



Draw the tangent RPR' meeting OQ , OQ' in R and R' .

Join PQ , and draw RW parallel to the axis, meeting PQ in W ;

Then, by the last Proposition,

$$PW = WQ.$$

And because RW is parallel to OP ,

$$\therefore OR : RQ :: PW : WQ.$$

But $PW = WQ$,

$$\therefore OR = RQ;$$

so $OR' = R'Q'$,

$$\therefore OR : RQ :: OR' : R'Q',$$

$$\therefore RR' \text{ is parallel to } QQ'.$$

Again, since PR is parallel to QV ,

$$\therefore OP : PV :: OR : RQ.$$

But $OR = RQ$,

$$\therefore OP = PV.$$

COR. From this it is manifest that if any number of parallel chords be drawn in a parabola, their middle points will all lie on the line parallel to the axis which passes through the point where the tangent drawn parallel to the chords meets the parabola.

DEF. Any line PV , drawn from a point P in the parabola parallel to the axis, is called a *Diameter*.


The point P is called the *Vertex* of the diameter PV ; and the tangent at P the *Tangent at the Vertex*.

The diameter consequently bisects all chords parallel to the tangent at the vertex, and the tangents at the extremities of any chord will intersect in the diameter corresponding to that chord.

DEF. A line QV , drawn parallel to the tangent at P from a point Q in the curve, is called the *Ordinate* to the diameter PV .

PROP. XV.

✓ 17. If QV be an ordinate to the diameter PV , then $QV^2 = 4 \cdot SP \cdot HV$.



Produce QV to meet the parabola in Q' ; and draw the tangents QO , $Q'O$, meeting VP produced in the point O . (*Prop. XIV.*)

Also let the tangent at P meet OQ in R , and join SP , SR , and SQ . Now since from the point R two tangents RP , RQ are drawn to the parabola, the triangle RPS is similar to the triangle RSQ , (*Prop. XII.*)

$\therefore$ the angle $SRP =$ the angle SQR .

But the angle $SQR =$ the angle STQ , (*Prop. VII.*)
 $=$ the angle POR ,

$\therefore$ the angle $SRP =$ the angle POR ,

and the angle $SPR =$ the angle OPR , (*Prop. V. Cor. 1*)

$\therefore$ the remaining angle $RSP =$ the remaining angle QRP ,

$\therefore$ the triangle SPR is similar to the triangle POR ,

$$\therefore SP : PR :: PR : PO,$$

$$\therefore PR^2 = SP \cdot PO,$$

$$= SP \cdot PV. \quad (\text{Prop. XIV.})$$

Again, since QV is parallel to PR ,


$$\therefore QV : PR :: OV : OP.$$

$$\text{But } OV = 2 \cdot OP, \quad (\text{Prop. XIV.})$$

CONIC SECTIONS.

PROP. XVII.

19. If two chords of a parabola intersect one another, the rectangles contained by their segments are in the ratio of the parameters of the diameters which bisect the chords.



Let the chords Qq , $Q'q'$ intersect one another in the point O .

Bisect Qq , $Q'q'$ in V and V' ; and draw the diameters PV , $P'V'$ parallel to the axis.

Also, through O draw OR parallel to $P'V'$; and through R draw RW parallel to QV .

Now, since Qq is divided equally in V and unequally in O ,

$$\begin{aligned} \therefore QO \cdot Oq &= QV^2 - OV^2, \quad (\text{Euclid, II. 5}) \\ &= QV^2 - RW^2, \\ &= 4SP \cdot PV - 4SP \cdot PW, \quad (\text{Prop. XV.}) \\ &= 4SP \cdot RO. \end{aligned}$$


So $Q'O \cdot Oq' = 4SP' \cdot RO$.

Hence $QO \cdot Oq : Q'O \cdot Oq' :: 4SP : 4SP'$.

By *Euclid*, II. 6, the same may be proved to be true if the point O be without the parabola.

PROP. XVIII. (omit)

20. If from an external point O a pair of tangents OQ , OQ' be drawn to the parabola, and the chord QQ' be joined, the area of the figure bounded by QQ' and the curve is two-thirds of the triangle QOQ' .



Draw the diameter OV meeting the curve in P ; and let the tangent at P meet OQ , OQ' in R and R' .

Join QP , $Q'P$; then

since $OR = RQ$,

$\therefore$ the triangle $OPR = \frac{1}{2}$ the triangle OPQ ,
 $= \frac{1}{2}$ the triangle VPQ .

So the triangle $OPR' = \frac{1}{2}$ the triangle VPQ ,

$\therefore$ the triangle $ORR' = \frac{1}{2}$ the triangle PQQ' .

Again, if through R and R' we draw the diameters Rp , $R'p'$; and at the points p and p' draw the tangents rpr , $r'p'r'$, we can prove in the same manner as before that

the triangle $Rrr = \frac{1}{2}$ the triangle QpP ,

and the triangle $R'r'r' = \frac{1}{2}$ the triangle $Q'p'P$.

Continuing in this manner to form new triangles by drawing diameters at the points r , r , and r' , r' , and tangents at the points where these diameters meet the curve, we can prove that the exterior triangles formed by the tangents are the halves of the interior triangles formed by joining the points of contact with the extremities of the chords.

And the same will hold however the number of the triangles be increased.

Hence the sum of all the exterior triangles will be equal to half the sum of all the interior triangles.

Now when the number of the triangles is increased indefinitely, the sum of the exterior triangles will represent the exterior figure $OQPQ'$, and the sum of the interior triangles the area of the interior figure $QPPQ'$. Hence

the area of the figure $OQPQ' = \frac{1}{2}$ the area of the figure $QPPQ'$,

$\therefore$ area of the figure $OQPQ' = \frac{1}{3}$ the area of triangle QOQ' ,

$\therefore$ area of the figure $QPPQ' = \frac{2}{3}$ the area of triangle QOQ' .

21. DEF. If with a point O on the normal at P as centre and OP as radius, a circle be described touching the parabola at P and cutting it in Q ; then when the point Q is made to approach indefinitely near to P , the circle is called the *Circle of Curvature* at the point P . (See fig. Prop. XIX.)

PROP. XIX.

The chord of the circle of curvature, at a point P of a parabola, drawn parallel to the axis $= 4SP$.


Let PT be the tangent, and PG the normal at the point P .

With centre O and radius OP describe a circle cutting the parabola in the point Q .

Draw RQX parallel to the axis meeting the circle in X and the tangent at P in R .

Also draw QV parallel to PR , and PW parallel to the axis; then

since RP touches the circle at P ,
 $\therefore RQ \cdot RX = PR^2$. (*Euclid*, III. 36.)



But $PR^2 = QV^2 = 4SP \cdot PV$, (*Prop. XV.*)

$$\therefore RQ \cdot RX = 4SP \cdot PV.$$

But $RQ = PV$,

$$\therefore RX = 4SP.$$

Now when the circle becomes the circle of curvature at P , the points R and Q move up to and coincide with P , and the lines RX and PV become equal.

Hence the chord of the circle of curvature parallel to the axis $= 4SP$.

COR. 1. If PU be the diameter of the circle of curvature, and PF the chord through the focus; then

since the angle $FPU =$ the angle WPU , (*Prop. VIII.*)

$$\therefore PF = PW = 4SP.$$

COR. 2. If SY be drawn at right angles to PT ; then the triangle PFU is similar to SYT ,

$$\therefore PU : PF :: SP : SY,$$

$$\text{or } PU : 4SP :: SP : SY.$$

PROP. XX.

If QVQ' be any ordinate to the diameter PI , the circle described through the three points P , Q , Q' will intersect the parabola in a fourth point, which depends only upon the position of P .


Draw the ordinate PN , and produce it to meet the parabola in P' ; then,

since the subtangent $= 2 \cdot AN$. (*Prop. VII.*)

The tangents at P and P' will meet the axis in the same point T .

Draw PR parallel to TP' , meeting the parabola in R , and QQ' in O ; then

$$PO \cdot OR : QO \cdot OQ' :: SP' : SP. \quad (\text{Prop. XVII.})$$



But $SP = ST = SP'$, (*Prop. VII.*)

$$\therefore PO \cdot OR = QO \cdot OQ'.$$

Hence by the converse of *Euclid III. Prop. 22*, the point R is on the circle which passes through P , Q , Q' .

COR. 1. Since TP and TP' are equally inclined to the axis, the lines QQ' , PR , which are parallel respectively to TP and TP' , are also equally inclined to the axis.

COR. 2. When the point V is brought indefinitely near to P , QQ' coincides with the tangent to the parabola at P , and becomes also a tangent to the circle at P , since Q and Q' are indefinitely near to each other. The circle therefore becomes the circle of curvature at the point P .

Hence if PR be drawn parallel to the tangent at P' , or be equally inclined to the axis with PT , it will meet the parabola in the point where the circle of curvature at P intersects the parabola.

PROBLEMS ON THE PARABOLA.

1. THE diameter of the circle described about the triangle BAC is equal to $5AS$. (See *fig. Prop. III.*)

2. If from the point G , GK be drawn at right angles to SP , then $PK = 2AS$. (See *fig. Prop. VII.*)

3. If the triangle SPG is equilateral, then SP is equal to the latus rectum. (See *fig. Prop. VII.*)

4. PQ is a common tangent to a parabola and the circle described on the latus rectum as diameter; prove that SP and SQ make equal angles with the latus rectum.

5. Prove that $PY \cdot PZ = SP^2$, and that $PY \cdot YZ = AS \cdot SP$. (See *fig. Prop. VII.*)

6. If PL be drawn at right angles to AP , meeting the axis in L , and PN be the ordinate of P , then $NL = 4AS$.

7. The tangent at any point P of a parabola meets the directrix and latus rectum produced in points equally distant from the focus.

8. Prove that $NY = TY$, and that $TP \cdot TY = TS \cdot TN$. (See *fig. Prop. VII.*)

9. If a circle be described about the triangle SPN , the tangent to it from $A = \frac{1}{2}PN$. (See *fig. Prop. VII.*)

10. If the ordinate of a point P bisect the subnormal of P' , the ordinate of P is equal to the normal of P' .

11. If from any point on the tangent to a parabola a line be drawn touching the parabola, the angle between this line and the line to the focus from the same point is constant.

12. A circle and parabola have the same vertex and axis. $BA'C$ is the double ordinate of the parabola which touches the circle at A' , the extremity of the diameter through the vertex A . PP' is any other ordinate of the parabola parallel to this, meeting the axis in N , and AB produced in R ; prove that the rectangle $RP \cdot RP'$ is proportional to the square of the tangent drawn from N to the circle.

13. Draw a parabola to touch a given circle at a given point, and such that its axis may touch the same circle in another given point.

14. If from the point of contact of a tangent to a parabola a chord be drawn, and another line be drawn parallel to the axis meeting the chord, tangent, and curve, this line will be divided by them in the same ratio as it divides the chord.

15. If the diameter PV meet the directrix in O , and the chord drawn through the focus parallel to the tangent at P in V ; prove that $PV = PO$.

16. Prove that the locus of the intersection of a diameter PV with the chord drawn through the focus parallel to the tangent at P is a parabola.

17. If a circle and parabola have a common tangent at P , and intersect in Q and R ; and QV , UR be drawn parallel to the axis of the parabola meeting the circle in V and U respectively, then VU is parallel to the tangent at P .

18. AB and AC are two lines at right angles to each other. From a fixed point C on AC , CR is drawn parallel to AB . On AR , produced if necessary, P is taken such that the perpendicular PN upon AB is equal to CR . Prove that the curve traced out by P is a parabola.

19. If from a point P of a circle PC be drawn to the centre; and R be the middle point of the chord PQ drawn parallel to a fixed diameter ACB , then the curve traced out by the intersection of CP and AR is a parabola.

20. If two equal tangents OQ , OQ' be cut by a third tangent, their alternate segments are equal.

21. E is the centre of the circle described about the triangle OQQ' . Prove that the circle described about the triangle QEQ' will pass through the focus. (See fig. Prop. XIII.)

22. PSp is any focal chord of a parabola. Prove that AP , Ap will meet the latus rectum in two points Q, q , whose distances from the focus are equal to the ordinates of p and P .

23. PSp is a focal chord of a parabola, lDr the directrix meeting the axis in D ; and Q any point on the curve. Prove that if QP, Qp be produced to meet the directrix in R, r , half the latus rectum is a mean proportional between DR, Dr .

24. OP and OQ are two tangents to a parabola. On QO produced, OQ' is taken equal to OQ ; prove that $OS \cdot PQ' = OP \cdot OQ$.

25. If QD be drawn at right angles to the diameter PV , then $QD^2 = 4AS \cdot PV$.

26. If through any point O on the axis of a parabola a chord POQ be drawn, and PM, QN be the ordinates of the points P and Q , prove that $AM \cdot AN = AO^2$.

27. If AP and AQ be drawn at right angles to each other from the vertex of a parabola, and PM, QN be the ordinates of P and Q , prove that the latus rectum is a mean proportional between AM and AN .

28. OAP is the sector of a circle whose centre is O . If the radius OA remain fixed while the angle AOP changes, the centre of the circle inscribed in the sector, AOP will trace out a parabola.

29. QSQ' is a focal chord parallel to AP ; $PN, QM, Q'M'$ are the ordinates of P, Q , and Q' . Prove that $SM^2 = AM \cdot AN$ and that $MM' = AP$.

30. PQ, PQ' are drawn from any point P cutting the ordinates $Q'V', QV$ in R' and R , prove that VR is to $V'R'$ in the triplicate ratio of QV to $Q'V'$.

31. On a chord of a parabola as diameter a circle is described cutting the parabola again in two points. If these points be joined, the portion of the axis between the two chords is equal to the latus rectum.

32. If OQ, OQ' be a pair of tangents to a parabola, and the chord QQ' be a normal to the curve at Q , then OQ is bisected by the directrix.

33. Two equal parabolas having the same focus and their axes in contrary directions intersect at right angles.

34. The radius of curvature at the extremity of the latus rectum is equal to twice the normal.

35. If from any point P of a parabola PF and PH be drawn making equal angles with the normal PG , then $SG^2 = SF \cdot SH$.

36. If a triangle be inscribed in a parabola, the points when the sides produced meet the tangents at the opposite angles are in the same straight line.

37. If the tangents OQ, OQ' be cut by a third tangent in R, R' , prove that

$$OR : RQ :: RQ' : OR'.$$

38. If from the vertex of a parabola chords be drawn at right angles to one another, and on them a rectangle be described, the curve traced out by the further angle is a parabola.

39. Prove that $2PY$ is a mean proportional between AP and the chord of the circle of curvature at the point P of the parabola drawn through the vertex A . (See fig. Prop. VII.)

40. If a circle described upon the chord of a parabola as diameter meet the directrix it also touches it, and all chords for which this is possible intersect in a point.

41. If a parabola roll upon another equal parabola, the vertices originally coinciding, the focus traces out the directrix.

42. The circle of curvature at the extremity of the latus rectum intersects the parabola on the diameter of curvature passing through the point of contact.

CHAPTER II.

THE ELLIPSE.

22. DEF. The *Ellipse* is the curve traced out by a point which moves in such a manner that its distance from a given fixed point continually bears the same ratio, *less than unity*, to its distance from a given fixed line. (See *Introduction*.)

PROP. I.

The focus and directrix of an ellipse being given, to find any number of points on the curve.

Let S be the focus and MX the directrix.

Draw SX at right angles to the directrix, and divide SX in the point A , so that SA may be to AX in the given fixed ratio less than unity; then

A is a point on the curve.

On XS produced take a point A' , such that


$$SA' : A'X :: SA : AX;$$

then A' will also be a point on the curve.

On the directrix take *any* point M ; and through M and S draw the line $MYSY'$, meeting AY and $A'Y'$, drawn at right angles to AA' , in the points Y and Y' .

On YY' as diameter describe a circle, and draw MPP' parallel to AA' , cutting the circle in the points P and P' ;

P and P' will be points on the ellipse.



Join PY , PY' , SP ; then since

$$SY : YM :: SA : AX, \quad (\text{Euclid, VI. 2})$$

$$\text{and } SY' : Y'M :: SA' : A'X, \quad (\text{Euclid, VI. 2})$$

$$\therefore SY : YM :: SY' : Y'M;$$

or, alternately,

$$SY : SY' :: YM : Y'M,$$

and the angle YPY' in a semicircle is a right angle,

$\therefore PY$ bisects the angle SPM ,*

$$\therefore SP : PM :: SY : YM,$$

$$:: SA : AX.$$

So we may show that

$$SP' : P'M :: SY : YM,$$

$$:: SA : AX,$$

$\therefore P$ and P' are points on the curve.

* For, if not, make the angle YPs equal to YPM ; then

$$sY : YM :: sP : PM. \quad (\text{Euclid, VI. 3.})$$

And since, if PY bisect sPM , PY' , being at right angles to PY , also bisects the angle sPM' ,

$$\therefore sY' : Y'M :: sP : PM. \quad (\text{Euclid, VI. A.})$$

$$\text{Hence } sY : YM :: sY' : Y'M,$$

$$\text{or } sY : sY' :: YM : Y'M,$$

$\therefore$ the points S and s coincide.

In the same way, by taking other points on the directrix, we may obtain as many more points on the curve as we please.

COR. 1. Since, corresponding to every point P on the curve, there is a point P' situated in precisely the same manner with respect to $A'Y'$ as P is with respect to AY , it is clear that if we make $A'S'$ equal to AS , and $A'X'$ equal to AX , and draw $X'M'$ at right angles to $A'X'$, the curve could be equally well described with S' as focus and $M'X'$ as directrix.

The ellipse is therefore symmetrical, not only with respect to the line AA' , but also with respect to the line OC drawn through the middle point of YY' at right angles to and bisecting AA' .

COR. 2. The line OP will bisect the angle SPS' .

Let OP meet SS' in G . Produce MP to meet $X'M'$ in M' , and draw OM' passing through the focus S' ; then

$$SP : PM :: S'P : PM',$$

or, alternately, $SP : S'P :: PM : PM'$. (1)

$$\text{Again, } SG : PM :: S'G : PM',$$

or, alternately, $SG : S'G :: PM : PM'$, (2)

$\therefore$ from (1) and (2)

$$SP : S'P :: SG : S'G,$$

$\therefore PG$ bisects the angle SPS' . (*Euclid*, VI. 3.)

It will be shown hereafter (*Prop.* XI.) that the normal to the ellipse at the point P also bisects the angle SPS' . Hence the ellipse and circle have the same tangent at the point P .

The ellipse will consequently touch all the infinite series of circles which can be described in the same manner as the one in the figure by taking different points on the directrix.

PROP. II.

23. If C be the middle point of AA' , then CA is a mean proportional between CS and CX ,

or $CS \cdot CX = CA^2$. (See fig. Prop. III.)

Since $SA' : A'X :: SA : AX$.

Alternately $SA' : SA :: A'X : AX$,

$\therefore SA' + SA : SA :: A'X + AX : AX$;

or $AA' : SA :: XX' : AX$,

$\therefore AA' : XX' :: SA : AX$.

or $CA : CX :: SA : AX$. (1)*

Again, $SA' : SA :: A'X : AX$,

$\therefore SA' - SA : SA :: A'X - AX : AX$;

or $SS' : SA :: AA' : AX$.

Alternately $SS' : AA' :: SA : AX$;

or $CS : CA :: SA : AX$. (2)

Hence from (1) and (2)

$CA : CX :: CS : CA$,

$\therefore CA^2 = CX \cdot CS$;

or CA is a mean proportional between CS and AX .

COR. Since the three lines CS , CA , CX are proportional, therefore, by the definition of duplicate ratio and *Euclid*, VI. 20 Cor.


$CA : CX :: CS^2 : CA^2$. (3)

PROP. III.

24. If P be any point on the ellipse, then

$SP + S'P = AA'$.

* N.B. The results (1), (2), (3), should be remembered, as they will frequently be referred to.



Since $SP : PM :: SA : AX$,
 and $A : AX :: A' : X'$, (*Prop. II.*)
 $\therefore SP : PM :: AA' : XX'$,
 So $S'P : PM' :: AA' : XX'$,
 $\therefore SP + S'P : PM + PM' :: AA' : XX'$.
 But $PM + PM' = MM' = XX'$,
 $\therefore SP + S'P = AA'$.

Cor. 1. By means of this property the ellipse may be practically described and the form of the curve determined.

Let a string, equal in length to AA' , have its ends fastened to two points S and S' ; and let it be kept stretched by means of the point of a pencil at P ; then since $SP + S'P$ will be always equal to AA' , the point P will trace out the ellipse.

Cor. 2. The line AA' is the longest line that can be drawn in the ellipse.

For, if any other line PQ be drawn, then

$$\begin{aligned} SP + SQ &> PQ, \\ \text{and } S'P + S'Q &> PQ, \\ \therefore SP + S'P + SQ + S'Q &> 2PQ, \\ \text{or } AA' &> PQ. \end{aligned}$$

25. **DEF.** If BCB' be drawn at right angles to ACA' , meeting the ellipse in B and B' , it will be seen further on (*Prop. XIII. Cor. 2*) that BCB' is the shortest chord that can be drawn through the *centre* of the ellipse. (*See fig. Prop. IV.*)

AA' is called the *Major Axis* and BB' the *Minor Axis* of the ellipse.

In most geometrical treatises the ellipse is defined as the curve traced out by a point which moves in such a manner that the sum of its distances from two fixed points is always the same; but it appears that the properties of the curve are more clearly exhibited by defining it in a manner analogous to the parabola, and deducing *immediately* from that definition the property in question.

Having now shown that one definition necessarily includes the other, we are at liberty in our future investigations to make use of whichever property is most convenient.


✎ PROP. IV.

26. If BC be the semi-minor axis of the ellipse, then

$$BC^2 = CA^2 - CS^2;$$

and if SL be the semi-latus rectum,

$$SL \cdot AC = BC^2.$$



Join SB , $S'B$; then

since $SB + S'B = AA'$, (*Prop. III.*)

and that $SB = S'B$,

$$\therefore SB = AC.$$

$$\text{But } BC^2 = SB^2 - CS^2,$$


$$\therefore BC^2 = CA^2 - CS^2.$$

COR. Conversely, a point will be inside or outside the ellipse according as the sum of its distances from the foci is less or greater than AA' .

28. DEF. If a point P' be taken on the ellipse near to P , (see fig. Prop. VI.) and PP' be joined, the line PP' produced, in the limiting position which it assumes when P' is made to approach indefinitely near to P , is called the *Tangent* to the ellipse at the point P .

PROP. VI.

If the tangent to the ellipse at any point P intersect the directrix in the point Z , and if S be the focus corresponding to the directrix on which Z is situated, then SZ will be at right angles to SP .



Let P' be a point on the ellipse near to P .

Draw the chord PP' , and produce it to meet the directrix in Z ; join SZ .

Draw PM , $P'M'$ at right angles to the directrix, and join SP , SP' .

Produce PS to meet the ellipse in the point Q ; then since the triangles $ZMP, ZM'P'$ are similar,

$$\begin{aligned}\therefore ZP : ZP' &:: MP : M'P', \\ &:: SP : SP',\end{aligned}$$

$\therefore SZ$ bisects the angle $P'SQ$. (*Euclid*, VI. Prop. A.)

Now when P' is indefinitely near to P , and PP' becomes the tangent at the point P , the angle PSI' becomes indefinitely small, while the angle QSI' approaches two right angles; and therefore the angle ZSP , being half of the angle $P'SQ$, becomes ultimately a right angle.

Hence when PZ becomes the tangent at the point P ,
the angle ZSP is a right angle,
or SZ is perpendicular to SP .

COR. 1. Conversely, if SZ be drawn at right angles to SP meeting the directrix in Z , and PZ be joined, PZ will be the tangent at P .

COR. 2. If ZP be produced to meet the other directrix in the point Z' , and $S'Z'$ be joined, then

$S'Z'$ is at right angles to $S'P$.

COR. 3. The tangents at the extremities of the latus rectum or double ordinate through the focus meet the axis produced in the point A .

PROP. VII.

The tangent to the ellipse at any point P makes equal angles with the focal distances SP and $S'P$.

Let the tangent at P meet the directrices in Z and Z' .


Draw MPM' at right angles to the directrices, meeting them in M and M' respectively; join $SZ, S'Z'$; then

$$SP : PM :: S'P : PM';$$

and since the triangles $MPZ, M'PZ'$, are similar,

$$PM : PZ :: PM' : PZ',$$

$$\therefore SP : PZ :: S'P : PZ'. \quad (\text{Ex æquali.})$$



Now in the triangles SPZ , $S'PZ'$, because the sides about the angles SPZ , $S'PZ'$ are proportional, and the angles PSZ , $PS'Z'$ are equal, being right angles, and the angles SZP , $S'Z'P$ are each less than a right angle,

$\therefore$ the triangles SPZ and $S'PZ'$ are similar, (*Euclid*, VI. 7)

$\therefore$ the angle $SPZ =$ the angle $S'PZ'$.

COR. If $S'P$ be produced to W ; then

the angle $SPZ =$ the angle WPZ .



PROP. VIII.

The tangents at the extremities of a focal chord intersect in the directrix.

Let PSQ be a focal chord, and let the tangent at P meet the directrix in Z .


Join SZ ; then

the angle ZSP is a right angle, (*Prop. VI.*)
 and $\therefore$ also the angle ZSQ is a right angle,
 $\therefore ZQ$ is the tangent at Q ; (*Prop. VI. Cor. 1*)
 or the tangents at the extremities of a focal chord intersect in the directrix.

PROP. IX.

29. If the tangent at P meet the axis major produced in T , and PN be the ordinate of the point P , then •

$$CT \cdot CN = CA^2.$$



Draw MPM' parallel to the axis major meeting the directrices in M and M' ; and produce $S'P$ to W ; then, since PT bisects the angle SPW , (*Prop. VII. Cor.*)

$$\therefore S'T : ST :: S'P : SP, \quad (\text{Euclid, VI. A.})$$

$$:: PM' : PM,$$

$$:: X'N : XN,$$

$$\therefore S'T + ST : S'T - ST :: X'N + XN : X'N - XN;$$

$$\text{or } 2CT : 2CS :: 2CX : 2CN,$$


$$\text{or } CT : CS :: CX : CN,$$

$$\therefore CT \cdot CN = CS \cdot CX,$$

$$= CA^2. \quad (\text{Prop. II.})$$

PROP. X.

If on the major axis of an ellipse as diameter a circle be described and a common ordinate NPQ be drawn meeting the ellipse in P and the circle in Q , then the tangents to the ellipse and circle respectively at the points P and Q will meet the major axis produced in the same point.)



Let the tangent to the ellipse at P meet the major axis produced in T ; join CQ , QT ; then, by the last Proposition,

$$CT \cdot CN = CA^2 = CQ^2,$$

$\therefore$ the angle CQT is a right angle.

And therefore QT is the tangent to the circle at Q ; ~~or~~ the tangents at P and Q meet the major axis produced in the same point T .


The circle described on AA' as diameter is called the *Auxiliary Circle* on account of the important aid that it affords in investigating the properties of the ellipse.

30. DEF. The line PG , drawn at right angles to the tangent PT , is called the *Normal* to the ellipse at the point P .

PROP. XI.

(If the normal at P meet the major axis in the point G ; then

$$SG : SP :: CS : CA.)$$



Since PG is at right angles to Tt ,

$\therefore$ the angle $GPT =$ the angle GPt .

But the angle $SPt =$ the angle $S'Pt$, (*Prop. VII.*)

$\therefore$ the angle $SPG =$ the angle $S'PG$,

or PG bisects the angle SPS' ,

$\therefore SG : S'G :: SP : S'P$, (*Euclid, VI. 3.*)

$\therefore SG : SG + S'G :: SP : SP + S'P$;

or $SG : SS' :: SP : AA'$;

or $SG : SP :: SS' : AA'$;

or $SG : SP :: CS : CA$.

Cor. Hence also,

$$S'G : S'P :: CS : CA.$$

Q. E. D. PROP. XII.

31. If the normal at P meet the major axis in G , and PN be the ordinate at the point P , then (*see fig. Prop. XI.*)

$$NG : NC :: BC^2 : AC^2.$$

Draw MPM' parallel to the axis meeting the directrices in M and M' ; join $SP, S'P$; then, since PG bisects the angle SPS' , (*Prop. XI.*)

$$\therefore S'G : SG :: S'P : SP,$$

$$:: PM' : PM,$$

$$:: X'N : XN,$$

$$\therefore S'G - SG : S'G + SG :: X'N - XN : X'N + XN;$$


$$\text{or } 2CG : SS' :: 2CN : XX'.$$

$$\begin{aligned}
 &\text{Alternately, } 2CG : 2CN :: SS' : XX'; \\
 &\quad \text{or } CG : CN :: CS : CX, \\
 &\qquad\qquad\qquad :: CS^2 : CA^2, \text{ (Prop. II. Cor.)} \\
 &\therefore CN - CG : CN :: CA^2 - CS^2 : CA^2; \\
 &\quad \text{or } NG : CN :: BC^2 : AC^2.
 \end{aligned}$$

PROP. XIII.

✓ 32. If PN be the ordinate of any point P on the ellipse ;
then

$$PN^2 : AN \cdot A'N :: BC^2 : AC^2.$$



Produce NP to meet the auxiliary circle in the point Q , and draw the tangents PT , QT meeting the major axis produced in the point T . (Prop. X.)

Join CQ , and let the normal at P meet the ellipse in G ; then, by the last Proposition,

$$NG : CN :: BC^2 : AC^2.$$

And rectangles of the same altitude are to another as their bases,

$$\therefore TN \cdot NG : TN \cdot CN :: BC^2 : AC^2;$$

$$\text{or } PN^2 : QN^2 :: BC^2 : AC^2. \text{ (Euclid, VI. 8, Cor.)}$$

$$\text{But } QN^2 = AN \cdot A'N,$$

since the angle AQA' in a semicircle is a right angle,

$$\therefore PN^2 : AN \cdot A'N :: BC^2 : AC^2.$$

COR. 1. Also $PN : QN :: BC : AC$.

This result is the basis of many of the future Propositions of the ellipse.

$$\begin{aligned} \text{But } CQ &= AC, \\ \therefore CR &= BC. \quad (1) \end{aligned}$$

Again, joining Rt ,


$$\begin{aligned} Ct : CU &:: PN : QN, \\ &:: CR : CQ, \\ \therefore Rt &\text{ is parallel to } QU, \\ \therefore CRt &\text{ is a right angle,} \\ \therefore Ct \cdot Cu &= CR^2. \quad (\text{Euclid, VI. 8, Cor}) \\ \text{But } CR &= BC, \quad (1) \\ \therefore Ct \cdot Cu &= BC^2. \end{aligned}$$

This proposition also admits of a demonstration similar to that given for the corresponding property of the hyperbola.

22

PROP. XV.

(33. If from the foci S and S' , SY and $S'Y'$ are drawn at right angles to the tangent at P , then Y and Y' are on the circumference of the auxiliary circle, and
 $SY \cdot S'Y' = BC^2$.)



Join SP , $S'P$, and produce SY and $S'P$ to meet in W ; and join CY ; then

since the angle $SPY =$ the angle WPY , (*Prop. VII. Cor.*)

and the angle $SY'P =$ the angle WYP ,

and the side PY is common to the triangles SPY , WPY ,

$\therefore$ the triangle $SPY =$ the triangle WPY in all respects,

$$\therefore SP = PW,$$

$$\therefore SP + S'P = S'W.$$

$$\text{But } SP + S'P = AA', \quad (\text{Prop. III.})$$

$$\therefore S'W = AA'.$$

Again $\because SC = S'C$ and $SY = YW$,

$$\therefore SC : S'C :: SY : YW,$$

$$\therefore CY \text{ is parallel to } S'W,$$

$$\therefore CY : S'W :: CS : SS',$$

$$\therefore CY = \frac{1}{2} S'W = CA.$$

$$\text{So } CY' = CA,$$

$\therefore Y$ and Y' are points on the auxiliary circle.

Next let YS be produced to meet the auxiliary circle in Z , and join ZY' ; then

since the angle ZYY' is a right angle,

$\therefore ZY'$ passes through the centre C ,

$\therefore$ the angle $SCZ =$ the angle $S'CY'$.

$$\therefore SZ = S'Y',$$

$$\therefore SY \cdot S'Y' = SY \cdot SZ,$$

$$= AS \cdot A'S, \quad (\text{Euclid, III. 35})$$

$$= CA^2 - CS^2, \quad (\text{Euclid, II. 5})$$


$$= BC^2. \quad (\text{Prop. IV.})$$

COR. If CD be drawn parallel to the tangent at T , meeting $S'P$ in E ; then

since the figure $CYPE$ is a parallelogram,
 $\therefore PE = CY = AC$.

PROP. XVI.

34. To draw a pair of tangents to an ellipse from an external point O .



With centre S' and radius equal to AA' describe a circle.

Join OS , OS' ; and let $S'O$ or $S'O$ produced meet the circle in the point I .

Now, if O be a point outside the circle MIM' , it is evident that OS is greater than OI ; and if O be inside the circle,

since $OS + OS' > AA'$ or $S'I$, (*Prop. V.*)

$\therefore OS > OI$.

With centre O and radius OS describe another circle cutting the former in the points M and M' , which it will always do since OS is greater than OI .

Join $S'M$, $S'M'$, meeting the ellipse in the points P and P' .

Join OP , OP' ; these will be the tangents required.

Join SP , SP' ; then since

$$SP + S'P = AA' = S'M,$$

$$\therefore SP = PM.$$

And $\because SP$, $PO = MP$, PO each to each,

$$\text{and } OS = OM,$$

$\therefore$ the angle $OPS =$ the angle OPM ,

$\therefore OP$ is the tangent at P . (*Prop. VII. Cor.*)

So OP' is the tangent at P' . •

PROP. XVII.

(If from a point O a pair of tangents OP , OP' be drawn to an ellipse, then OP and OP' will subtend equal angles at either focus.)

Join SP , $S'P$; SP' , $S'P'$; and produce $S'P$, $S'P'$ to M and M' , making PM equal to SP , and $P'M'$ equal to SP' .

Join OM , OM' ; OS , OS' .

Then since OP , $PS = OP$, PM , each to each,

and the angle $OPS =$ the angle OPM , (*Prop. VII. Cor.*)

$$\therefore OS = OM,$$

and the angle $OSP =$ the angle OMP .

$$\text{So } OS = OM',$$

and the angle $OSP' =$ the angle $OM'P'$,

$$\therefore OM = OM'.$$

Again, $\because S'M = S'P + SP = AA'$,

and $S'M' = S'P' + SP' = AA'$.

$$\therefore S'M = S'M'.$$

And $\because OS'$, $S'M = OS'$, $S'M'$, each to each,

$$\text{and } OM = OM',$$

$\therefore$ the angle $OS'M =$ the angle $OS'M'$,

and the angle $OMS' =$ the angle $OM'S'$.

Then Tq and $T'q'$ will be tangents to the auxiliary circle. (*Prop. X.*)

Let Tq and $T'q'$ be produced to meet in o ; join Co meeting the chord qq' in v , and the circle in p .

Now, since the corresponding ordinates of the ellipse and auxiliary circle are in the constant ratio of BC to AC , the three lines ol , pm , vn drawn at right angles to AA' will pass through the points O , P , V respectively.

For, according as O is the point where ol meets TQ or $T'Q'$ we shall have

$$\begin{aligned} lO : lo &:: NQ : Nq, \\ &:: BC : AC; \\ \text{or } lO : lo &:: N'Q' : N'q', \\ &:: BC : AC, \end{aligned}$$

$\therefore Oo$ is perpendicular to AA' .

So Pp and Vv are perpendicular to AA' ,

$\therefore Oo, Pp, Vv$ are parallel.

Hence (1.) $QV : VQ' :: qv : vq'$.

But $qv = vq'$ from the circle,

$\therefore QV = VQ'$;

or QQ' is bisected in V .

(2.) Since $NQ : Nq :: N'Q' : N'q'$

it is evident that QQ' and qq' will meet the axis produced in the same point.

Also the tangents to the ellipse and circle at P and p respectively will meet the axis in the same point.

Now in the circle the tangent at p is manifestly parallel to qq' ,

and $NQ : Nq :: mP : mp$,

$\therefore$ the tangent at P is parallel to QQ' .

(3.) If Cq be joined, since the angle Cqo is a right angle, and Co is perpendicular to qq' ,

$\therefore Cv : Cq :: Cq : Co$, (*Euclid, VI. 8 Cor.*)

or, since $Cq = Cp$,

$Cv : Cp :: Cp : Co$.

E

$$\begin{aligned}
 &\text{But } Cv : Cp :: CV : CP, \\
 &\text{and } Cp : Co :: CP : CO, \\
 &\therefore CV : CP :: CP : CO, \\
 &\therefore CO \cdot CV = CP^2.
 \end{aligned}$$

COR. From this it is manifest that if any number of chords be drawn parallel to each other in an ellipse, their middle points will all lie on the line drawn from the centre to the point where the tangent parallel to the chord meets the ellipse.


DEF. The line PCP' drawn through the centre of an ellipse and meeting the curve in P and P' , is called a *Diameter*.

The diameter consequently bisects all chords parallel to the tangents at its extremities; and the tangents at the extremities of any chord will intersect the diameter corresponding to that chord in the same point. Δ

36. DEF. If CD be drawn parallel to the tangent at P , then CD is said to be *conjugate* to CP .

PROP. XIX.

(In the ellipse if CD be conjugate to CP , then will CP be conjugate to CD .)



Draw the ordinates PN , DR , and produce them to meet the auxiliary circle in the points p , d .

Join CP , Cp ; CD , Cd ; and draw the tangents TP , Tp ; $T'D$, $T'd$.

Now, since CD is parallel to PT ,

$\therefore$ the triangle PNT is similar to the triangle $DR C$.

$$\therefore TN : CR :: PN : DR,$$

$$:: Np : Rd, \quad (\text{Prop. XIII. Cor.})$$

$\therefore Tp$ is parallel to Cd ,

$\therefore$ the angle pCd is a right angle,

$\therefore Cp$ is parallel to $T'd$,

$\therefore$ the triangle pCN is similar to the triangle $dT'R$,

$$\therefore NC : RT' :: Np : Rd,$$

$$:: NP : RD,$$

$\therefore CP$ is parallel to DT' ,

$\therefore CP$ is conjugate to CD .

COR. Since CRd and CNp are each similar to dRT' ,
(*Euclid*, VI. 8)

$\therefore$ the triangle CRd is similar to the triangle CNp ,

and the side Cd = the side Cp ,

$\therefore$ the triangle CRd = the triangle CNp in all respects,

$$\therefore CN = Rd, \text{ and } CR = Np.$$

$$\text{Hence } DR : CN :: DR : Rd,$$

$$:: BC : AC;$$

$$\text{also } PN : CR :: PN : Np,$$

$$:: BC : AC.$$


PROP. XX.

(37. If CP and CD be conjugate semi-diameters, and PN , DR be the ordinates of the points P and D ; then

$$(1.) \quad CN^2 + CR^2 = AC^2.$$

$$(2.) \quad PN^2 + DR^2 = BC^2.$$

$$(3.) \quad CP^2 + CD^2 = AC^2 + BC^2. \quad \rangle$$



Produce NP , Rd to meet the auxiliary circle in the points p , d ; then

$$\begin{aligned} CN &= Rd, \quad (\text{Prop. XIX. Cor.}) \\ \therefore CN^2 + CR^2 &= Rd^2 + CR^2, \\ &= Cd^2, \\ &= CA^2. \end{aligned}$$

Again, $PN : Np :: BC : AC$,

$$\therefore PN^2 : Np^2 :: BC^2 : AC^2.$$

So $DR^2 : Rd^2 :: BC^2 : AC^2$,

$$\therefore PN^2 + DR^2 : Np^2 + Rd^2 :: BC^2 : AC^2;$$

$$\begin{aligned} \text{but } Np^2 + Rd^2 &= CR^2 + CN^2, \\ &= AC^2, \end{aligned}$$

$$\therefore PN^2 + DR^2 = BC^2,$$

$$\text{and } CN^2 + CR^2 = AC^2,$$

$$\therefore CP^2 + CD^2 = AC^2 + BC^2.$$

38. DEF. A line QV drawn parallel to the tangent at P , and meeting CP in V , is called an *Ordinate* to the diameter CP .

PROP. XXI.

(If QV be any ordinate to the diameter PCP' , and CD be conjugate to CP ; then


$$QV^2 : PV \cdot P'V :: CD^2 : CP^2.)$$

Draw the tangent UQW meeting CP and CD produced in U and W ; and draw QR parallel to CP , meeting CD in R .

Now, since $CR : CD :: CD : CW$, (*Prop. XVIII.*)

$\therefore CR^2 : CD^2 :: CR : CW$, (*Euclid, VI. 20 Cor.*)

or $QV^2 : CD^2 :: UV : CU$.



Again,

since $CU : CP :: CP : CV$, (*Prop. XVIII.*)

$\therefore CU : CV :: CP^2 : CV^2$, (*Euclid, VI. 20, Cor.*)

$\therefore CU - CV : CU :: CP^2 - CV^2 : CP^2$,

or $UV : CU :: PV \cdot P'V : CP^2$.

Hence $QV^2 : CD^2 :: PV \cdot P'V : CP^2$,


or $QV^2 : PV \cdot P'V :: CD^2 : CP^2$.

() PROP. XXII.

(39. The area of any parallelogram formed by drawing tangents to an ellipse at the extremities of a pair of conjugate

diameters is equal to the rectangle contained by the axes of the ellipse.)

Let PCP' , DCD' be a pair of conjugate diameters, and let a parallelogram be formed by drawing tangents at the points P , P' , D , D' .



Let the tangent at P meet CA produced in T ; join $D'T$.

Draw the ordinates PN , DR , $D'R'$; then since PT is parallel to CD' , the parallelogram PD' is double the triangle CTD' , and therefore equal to the rectangle contained by CT and $D'R'$.

Now $D'R' : CN :: BC : AC$, (*Prop. XIX. Cor.*)

$\therefore CT \cdot D'R' : CT \cdot CN :: BC : AC$, (*Euclid, VI. 1*)

$:: BC \cdot AC : AC^2$. (*Euclid, VI. 1*)

But $CT \cdot CN = AC^2$,

$\therefore CT \cdot D'R' = AC \cdot BC$,

$\therefore$ the parallelogram $LL' = 4$ the parallelogram PD' ,

$= 4AC \cdot BC$,

$= AA' \cdot BB'$.


COR. If PF be drawn at right angles to DCD' meeting CD' in F ; then

$$PF \cdot CD' = \text{area of parallelogram } PD', \\ = AC \cdot BC.$$

PROP. XXIII.

(40. If CP and CD be conjugate diameters, and PF be drawn at right angles to CD meeting CA in G , then

$$PF \cdot PG = BC^2. \quad \bullet$$



Draw the ordinate PN , and produce it to meet CD' in K .

Also draw Pn at right angles to CB , and let the tangent at P meet CB produced in t .


Now, since the angles at N and F are right angles, it is evident that a circle may be described about the quadrilateral figure $NKFG$;

$$\begin{aligned} \therefore PG \cdot PF &= PN \cdot PK, \quad (\text{Euclid, III. 36 Cor.}) \\ &= Ct \cdot Cn, \\ &= BC^2. \quad (\text{Prop. XIV.}) \end{aligned}$$

} PROP. XXIV.

(41. If P be any point on the ellipse, and CD be conjugate to CP , then

$$SP \cdot S'P = CD^2.)$$



Draw the normal PG and produce it to meet CD' in F ; then since CD' is parallel to the tangent at P ,

$\therefore PF$ is at right angles to CD' ,

$$\therefore PF \cdot CD = AC \cdot BC, \quad (\text{Prop. XXII. Cor.})$$

$$\text{and } PF \cdot PG = BC^2 = BC \cdot BC, \quad (\text{Prop. XXIII.})$$

$$\therefore CD : PG :: AC : BC. \quad (1)$$

$$\text{Again, } SP : SG :: CA : CS, \quad (\text{Prop. XI.})$$

$$S'P : S'G :: CA : CS. \quad (\text{Prop. XI.})$$

$$\text{Compounding } SP \cdot S'P : SG \cdot S'G :: CA^2 : CS^2,$$

$$\therefore SP \cdot S'P : SP \cdot S'P - SG \cdot S'G :: CA^2 : CA^2 - CS^2. \times$$

$$\text{But } SP \cdot S'P - SG \cdot S'G = PG^2, \quad (\text{Euclid, VI. Prop. B})$$

$$\therefore SP \cdot S'P : PG^2 :: CA^2 : BC^2. ($$


$$\text{But from (1) } CD^2 : PG^2 :: CA^2 : BC^2,$$

$$\therefore SP \cdot S'P = CD^2.$$

This proposition may also be very easily deduced from *Prop. XV.*

PROP. XXV.

42. The area of the ellipse is to the area of the auxiliary circle as BC to AC .)



Let PN and $P'N'$ be two ordinates of the ellipse near together.

Produce NP , $N'P'$, to meet the auxiliary circle in Q and Q' .

Draw Pm , Qn , perpendicular to $Q'N'$.

Then

the parallelogram PN' : the parallelogram QN' :: $PN : QN$,
:: $BC : AC$.

And the same will be true for all the parallelograms that can be similarly described in the ellipse and auxiliary circle.

Hence the sum of all the parallelograms inscribed in the ellipse is to the sum of all the parallelograms inscribed in the circle as BC to AC .

And this holds however the number of parallelograms be increased.

But when the number of parallelograms is increased, and the breadth of each diminished indefinitely, the sum of the parallelograms inscribed in the ellipse will be equal to the area of the ellipse, and the sum of those inscribed in the circle to the area of the circle. Hence


the area of the ellipse : the area of the circle :: $BC : AC$.

43. DEF. If with a point O on the normal at P as centre, and OP as radius, a circle be described touching the ellipse at P , and cutting it in Q ; then, when the point Q is made to approach indefinitely near to P , the circle is called the *Circle of Curvature* at the point P .

PROP. XXVI.

(If PH be the chord of the circle of curvature at the point P of an ellipse, which passes through the centre; then
 $PH \cdot CP = 2CD^2$.)

Let PT be the tangent, and PG the normal at the point P .



With centre O , and radius OP , describe a circle cutting the ellipse in the point Q .

Draw RQW parallel to CP , meeting the circle in W , and TP produced in R .

Also draw QV parallel to PR , meeting the diameter PP' in V ; then since RP touches the circle at P ,

$$\therefore RQ \cdot RW = PR^2, \text{ (Euclid, III. 36)}$$

$$\text{or } PV \cdot RW = QV^2.$$

But $QV^2 : PV \cdot P'V :: CD^2 : CP^2$, (*Prop. XXI.*)

$$\therefore PV \cdot RW : PV \cdot P'V :: CD^2 : CP^2,$$

$$\text{or } RW : P'V :: CD^2 : CP^2.$$

Now, when the circle becomes the circle of curvature at P , the points R and Q move up to, and coincide with P , and the lines RW and PH become equal, while

$P'V$ becomes equal to PP' , or $2CP$.

Hence, $PH : 2CP :: CD^2 : CP^2$,

$$\therefore PH \cdot CP : 2CP^2 :: 2CD^2 : 2CP^2,$$

$$\therefore PH \cdot CP = 2CD^2.$$

PROP. XXVII.

(If PU be the diameter of the circle of curvature at the point P of the ellipse, and PF be drawn at right angles to CD ; then

$$PU \cdot PF = 2CD^2.)$$

Since the triangle PHU is similar to the triangle $PF C$,

$$\therefore PU : PH :: CP : PF,$$

$$\therefore PU \cdot PF = PH \cdot CP,$$

$$= 2CD^2. \text{ (Prop. XXVI.)}$$

PROP. XXVIII.

(If PI be the chord of the circle of curvature through the focus of the ellipse; then

$$PI \cdot AC = 2CD^2.)$$

Let PI meet CD in E ; then, since the triangles PIU and PEF are similar,

$$\therefore PI : PU :: PF : PE.$$

But $PE = AC$, (*Prop. XV. Cor.*)

$$\therefore PI : PU :: PF : AC,$$

$$\begin{aligned}\therefore PI \cdot AC &= PU \cdot PF, \\ &= 2CD^2. \text{ (*Prop. XXVII.*)}\end{aligned}$$

PROP. XXIX.


(44. If two chords of an ellipse intersect one another, the rectangles contained by their segments are proportional to the squares of the diameters parallel to them.)

Let POP' be any chord drawn through the point O , and let CD be the semi-diameter parallel to it

Draw the ordinates NP , $N'P'$, MD . and produce them to meet the auxiliary circle in Q , Q' , D' ; then

since $NP : NQ :: N'P' : N'Q'$, (*Prop. XIII. Cor.*)

it is evident that PP' and QQ' will meet the axis produced in the same point T .



Also since $NP : NQ :: MD : MD'$, (*Prop. XIII. Cor.*)

and TPP' is parallel to CD ,

$\therefore TQQ'$ is parallel to CD' .

Draw EO parallel to NQ or $N'Q'$, and produce it to meet QQ' in O' ; then

$$* \quad PO : QO' :: TO : TO',$$

$$\text{and } P'O : Q'O' :: TO : TO',$$

$$\begin{aligned} \therefore PO \cdot P'O : QO' \cdot Q'O' &:: TO^2 : TO'^2, \\ &:: CD^2 : CD'^2, \\ &:: CD^2 : AC^2. \end{aligned}$$

Alternately, $PO \cdot P'O : CD^2 :: QO' \cdot Q'O' : AC^2$.

Again, if through the point O any other chord pOp' be drawn,

$$\text{since } EO : EO' :: BC : AC,$$

it is manifest that the corresponding chord qq' in the auxiliary circle will pass through the point O' ; and if Cd be the semi-diameter parallel to pp' we shall have as before,

$$pO \cdot p'O : Cd^2 :: qO' \cdot q'O' : AC^2.$$

But $QO' \cdot Q'O' = qO' \cdot q'O'$, (*Euclid*, III. 35)

$$\therefore PO \cdot P'O : CD^2 :: pO \cdot p'O : Cd^2,$$

$$\text{or } PO \cdot P'O : pO \cdot p'O : CD^2 : Cd^2.$$

The same result may be shown to be true when the point O is without the ellipse.


PROP. XXX. ✓

(If QVQ' be any ordinate to the diameter CP , the circle described through the three points P, Q, Q' will intersect the ellipse in a fourth point, which depends only upon the position of P .)

Draw the ordinate PN , and produce it to meet the ellipse in P' ; then, since, if NT be the subtangent of either P or P' ,

$$CT \cdot CN = AC^2, \text{ (Prop. IX.)}$$

therefore the tangents at P and P' will meet the major axis produced in the same point T .



Draw PR parallel to TP' , meeting the ellipse in R , and QQ' in O ; then if CD and CD' be drawn parallel respectively to TP and TP' , meeting the ellipse in D and D' ,

$$PO \cdot OR : QO \cdot OQ' :: CD'^2 : CD^2. \quad (\text{Prop. XXIX.})$$

But $CD' = CD$, since $CP' = CP$,

$$\therefore PO \cdot OR = QO \cdot OQ'.$$

Hence, by the converse of *Euclid* III. *Prop.* 22, the point R is on the circle which passes through P , Q , Q' .

COR. When the point V is brought indefinitely near to P , QQ' coincides with the tangent to the ellipse at P , and becomes also a tangent to the circle at P since Q and Q' are indefinitely near to each other. The circle therefore becomes the circle of curvature at the point P .

Hence, if PR be drawn parallel to the tangent at P' , or be equally inclined to the axis with PT , it will meet the ellipse in the point where the circle of curvature at P intersects the ellipse.

PROBLEMS ON THE ELLIPSE.

1. In what position of P is the angle SPS' greatest?
2. The latus rectum is a third proportional to the axis major and axis minor.
3. Construct on the axis minor as base, a rectangle which shall be to the triangle SLS' in the duplicate ratio of the major axis to the minor axis, L being the extremity of the latus rectum.
4. If a series of ellipses be described having the same major axis; the tangents at the extremities of their latera recta will all meet the minor axis in the same point.
5. Find the locus of the centres of all the ellipses having the same focus, and their major axes of the same length, and touching a given straight line.
6. Given the foci, it is required to describe an ellipse touching a given straight line.
7. If PT be a tangent to an ellipse, meeting the axis in T , and $AP, A'P$, be produced to meet the perpendicular to the major axis through T in Q and Q' , then $QT = Q'T$.
8. If the angle SBS' be a right angle, prove that $CA^2 = 2CB^2$.
9. If CP be a semi-diameter, and AQO be drawn parallel to CP meeting the curve in Q , and CB produced in O , then $CP^2 = AO \cdot AQ$.
10. If AB, CD , which are not parallel, make equal angles with either axis, the lines AC, BD , as also AD, BC , will make equal angles with either axis.

11. PSp is any focal chord. PA and pA are produced to meet the directrix in Q and q . Prove that the angle $Q Sq$ is a right angle.

12. If a circle be described touching the axis major in one focus, and passing through one extremity of the axis minor; AC will be a mean proportional between the diameter of this circle and BC .

13. If $PQQ'P'$ be a chord of the auxiliary circle, and a circle be described on the minor axis as diameter, cutting the chord in Q and Q' , then $PQ \cdot P'Q = CS^2$.

14. If PG be the normal at P , and GL be drawn at right angles to SP , then $PL = \frac{1}{2}$ latus rectum.

15. The sum of the squares of the normals at the extremities of conjugate diameters is constant.

16. If on the normal at P , PQ be taken equal to the semi-conjugate diameter CD , the locus of Q is a circle whose radius is $AC - BC$.

17. Find the locus of the intersection of a pair of tangents at right angles to each other.

18. P is any point on an ellipse. To any point Q on the curve draw AQ , $A'Q$, meeting NP in R and S , and prove that $NR \cdot NS = NP^2$.

19. If PG be a normal, and GL perpendicular to SP , the ratio of GL to PN is constant.

20. If NP produced meet the tangent at the extremity of the latus rectum in Q , then $QN = PS$.

21. In an ellipse the tangent at any point makes a greater angle with the focal distance than with the perpendicular on the directrix.

22. A diameter of an ellipse, parallel to the tangent at any point, meets the focal distances of the point, and from the points of intersection lines are drawn perpendicular to the focal distances. Prove that these lines intersect in the axis minor.

23. The subnormal is a third proportional to CT and BC .

24. If PN be the ordinate of P , prove that $NY : NY' :: PY : PY'$. (See fig. Prop. XV.)

25. If from C lines be drawn parallel and perpendicular to the tangent at P , they inclose a part of one of the focal distances of that point equal to the other.

26. If P be a fixed point on an ellipse, and QQ' an ordinate to CP , the circle QPQ' will meet the ellipse in a fixed point.

27. P is any point on an ellipse. Draw PP' parallel to the axis major, and through P' draw $P'Q, P'Q'$, making equal angles with the major axis. Join QQ' ; then QQ' is parallel to the tangent at P .

28. What parallelogram circumscribing an ellipse has the least area?

29. When is the square of the sum of conjugate diameters least?

30. Given the axes of an ellipse, and the position of one focus, and of one point in the curve, give a geometrical construction for finding the centre.

31. If lines drawn through any point of an ellipse to the extremities of any diameter meet the conjugate CD in M and N , then $CM \cdot CN = CD^2$.

32. If CP and CD be conjugate, prove that

$$(SP - AC)^2 + (SD - AC)^2 = SC^2.$$

33. If CP and CD be conjugate, and BP, BD be joined, as also AD, AP , these latter meeting in O , then $BDOP$ is a parallelogram. When is the area greatest?

34. If PSp, QCq be two parallel chords through the focus and centre of an ellipse, prove that

$$SP \cdot Sp : CQ \cdot Cq :: BC^2 : AC^2.$$

35. If the tangent at the vertex A cut any two conjugate diameters in T and t , then $AT \cdot At = AC^2$. ~~BC^2~~ AC^2 .

36. If the tangents at three points P, Q, R intersect in R', Q', P' , prove that

$$PR' \cdot P'Q \cdot Q'R = PQ' \cdot R'Q \cdot P'R.$$

37. If a circle be described touching SP , $S'P$ produced, and the major axis of the ellipse, find the locus of the centre.

38. If from the extremities of the axes of an ellipse any four parallel lines be drawn, the points in which they cut the curve are the extremities of conjugate diameters.

39. If two equal and similar ellipses have a common centre, the points of intersection are at the extremities of diameters at right angles to one another.

40. If PSQ be a focal chord, and X the foot of the directrix, XP and XQ are equally inclined to the axis.

41. OP , OQ are tangents to an ellipse, and PQ is produced to meet the directrices in R , R' , prove that

$$RP \cdot R'P : RQ \cdot R'Q :: OP^2 : OQ^2.$$

42. NPQ is a common ordinate to the ellipse and auxiliary circle. PR , QR are normals at P and Q intersecting in R . The locus of R is a circle whose radius is $AC + BC$.

43. If the conjugate to CP meet SP , $S'P$, or these produced in E , E' ; then $SE = S'E'$, and the circles circumscribing SCE , $S'CE'$ are equal.

44. The locus of the middle points of all focal chords in an ellipse is a similar ellipse.

45. The circle described about the triangle SBS' will cut the minor axis in the centre of the circle of curvature at B .

46. The locus of the centre of the circle inscribed in the triangle SPS' is an ellipse.

47. If a circle be described intersecting an ellipse in four points, and chords be drawn through the points of intersection, diameters parallel to the chords will be equal.

48. An ellipse slides between two lines at right angles to each other, find the locus of its centre.

49. If from the focus S perpendiculars be drawn upon the conjugate diameters CP , CD , these perpendiculars produced backward will intersect CD and CP in the directrix.

50. Find the point at which the radius of curvature is a mean proportional between the major and minor axes.

51. The circle of curvature at a point, where the conjugate diameters are equal, meets the ellipse again at the extremity of the diameter.

52. The locus of the intersection of lines drawn from A, A' at right angles to $AP, A'P$ is an ellipse.

53. If a quadrilateral figure be inscribable in two ellipses whose major axes are parallel or perpendicular, any two of its opposite angles will be equal to two right angles.

54. If CN, NP are the abscissa and ordinate of a point P on a circle whose centre is C , and NQ be taken equal to NP , and be inclined to it at a constant angle, the locus of Q is an ellipse.

55. If two ellipses having the same major axes can be inscribed in a parallelogram, the foci will be on the corners of an equiangular parallelogram.

56. Two ellipses, whose major axes are equal, have a common focus. Prove that they intersect in two points only.

57. A circle described about the triangle SPS' cuts the minor axis in R on the opposite side to P . Prove that SR varies as the normal PG .

58. If r and R be the radii of the circles inscribed in and about the triangle SPS' , prove that $R \cdot r$ varies as $SP \cdot S'P$.

59. The circle described upon PG as diameter cuts $SP, S'P$ in K and L . Prove that KL is bisected by PG , and is perpendicular to it.

60. If from S' a line be drawn parallel to SP , it will meet SY in the circumference of a circle.

61. T and t are the points where the tangent at P meets the axes. CP is produced to meet in L the circle described about the triangle TCt ; prove that PL is half the chord of the circle of curvature at P in the direction of C , and that $CP \cdot CL$ is constant.

62. About the triangle PQR an ellipse is described, having its centre at the point where the lines drawn from P, Q, R to the middle points of the opposite sides meet. CP, CQ, CR are produced to meet the ellipse in P', Q', R' . Prove that

the tangents at P' , Q' , R' form a triangle similar to PQR , and four times as large.

63. Lines from Y and Y' perpendicular to the major axis cut the circles on SP , $S'P$ as diameters in I and J . Prove that IS , and JS' , when produced, intersect BC in the same point.

64. If from the ends of any diameter chords be drawn to any point in the ellipse, the diameters parallel to these chords will be conjugate.

65. If T be the angle between tangents at the extremities of a focal chord, and O the angle subtended by the chord at the other focus, then

$$2T + O = 2 \text{ right angles.}$$

66. The acute angles which SP , SQ make with the tangents are complementary. Prove that BC^2 is a mean proportional between the areas of the triangles SPS' , SQS' . Also, show that the problem is impossible unless $BC < CS$.

67. A series of ellipses have their equal conjugate diameters of the same magnitude. One of these diameters is fixed and common, while the other varies. The tangents drawn from any point in the fixed diameter produced will touch the ellipses in points situated on a circle.

68. If on the longer side of a rectangle as major axis an ellipse be described, passing through the intersection of the diagonals, and lines be drawn from any point of the ellipse exterior to the rectangle to the ends of the remote side, they will divide the major axis into segments, which are in geometric progression.

69. From any point P of an ellipse PQ is drawn at right angles to SP meeting the diameter conjugate to CP in Q . Prove that PQ varies inversely as the perpendicular from P on the major axis.

70. In an ellipse SQ and $S'Q$, drawn at right angles to a pair of conjugate diameters, intersect in Q . Prove that the locus of Q is a concentric ellipse.

CHAPTER III.

THE HYPERBOLA.

45. DEF. The *Hyperbola* is the curve traced out by a point which moves in such a manner that its distance from a given fixed point continually bears the same ratio, *greater than unity*, to its distance from a given fixed line. (*See Introduction.*)

PROP. I.

(The focus and directrix of a hyperbola being given, to find any number of points on the curve.)

Let S be the focus, and MX the directrix.

Draw SX at right angles to the directrix, and divide SX in the point A , so that SA may be to AX in the given fixed ratio, greater than unity; then

A is a point on the curve.

On SX produced take a point A' , such that


$$SA' : A'X :: SA : AX;$$

then A' will also be a point on the curve.

On the directrix take *any* point M ; and through S and M draw the line $SYM Y'$, meeting AY and $A'Y'$, drawn at right angles to AA' , in the points Y and Y' ;

On YY' as diameter describe a circle, and draw PMP' parallel to AA' , cutting the circle in the points P and P' ;

P and P' will be points on the hyperbola.



Join PY, PY', SP ; then since

$$SY : YM :: SA : AX, \quad (\text{Euclid, VI. 2.})$$

$$\text{and } SY' : Y'M :: SA' : A'X, \quad (\text{Euclid, VI. 2.})$$

$$\therefore SY : YM :: SY' : Y'M;$$

or, alternately, $SY : SY' :: YM : Y'M$,

and the angle YPY' in a semicircle is a right angle,

$$\therefore PY \text{ bisects the angle } SPM,*$$

$$\therefore SP : PM :: SY : YM,$$

$$:: SA : AX.$$

So we may show that

$$SP' : P'M :: SY' : Y'M,$$

* For, if not, make the angle YPm equal to YPS ; then

$$SY : Ym :: SP : Pm, \quad (\text{Euclid, VI. 3})$$

and, since, if PY bisect SPm , PY' being at right angles to PY , also bisects the angle between MP and SP produced;

$$\therefore SY' : Y'm :: SP : Pm, \quad (\text{Euclid, VI. A.})$$

$$\text{Hence } SY : Ym :: SY' : Y'm,$$

$$\text{or } SY : SY' :: Ym : Y'm;$$

$\therefore$ the points M and m coincide.

$$:: SA : AX,$$

$\therefore P$ and P' are points on the curve.

In the same way, by taking other points on the directrix, we may obtain as many more points on the curve as we please.

COR. 1. Since, corresponding to every point P on the curve, there is a point P' situated in precisely the same manner with respect to $A'Y'$ as P is with respect to AY , it is clear that if we make $A'S'$ equal to AS , and $A'X'$ equal to AX , and draw $X'M'$ at right angles to AX' , the curve could be equally well described with S' as focus and $M'X'$ as directrix.

The hyperbola is therefore symmetrical, not only with respect to the line AA' , but also with respect to the line OC' drawn through the middle point of YY' at right angles to and bisecting AA' .

COR. 2. The line OP produced will bisect the angle SPW' between SP and $S'P$ produced.

Produce OP and $S'S$ to meet in G . Produce PM to meet $X'M'$ in M' , and draw OS' passing through the point M' ; then

$$SP : PM :: S'P : PM',$$

$$\text{or, alternately, } SP : S'P :: PM : PM'. \quad (1.)$$

$$\text{Again, } SG : PM :: S'G : PM',$$

$$\text{or, alternately, } SG : S'G :: PM : PM'. \quad (2.)$$

$$\therefore \text{ from (1) and (2)}$$

$$SP : S'P :: SG : S'G.$$

$$\therefore PG \text{ bisects the angle } SPW. \quad (\text{Euclid, VI. A.})$$

It will be shown hereafter (*Prop. IX.*) that the normal to the hyperbola at the point P also bisects the angle SPW . Hence the hyperbola and circle have the same tangent at the point P . The hyperbola will consequently touch all the infinite series of circles which can be described in the same manner as the one in the figure, by taking different points on the directrix.

PROP. II.

(46. If C be the middle point of AA' ; then CA is a mean proportional between CS and CX ,)

or $CS \cdot CX = CA^2$. (See fig. Prop. III.)

Since $SA' : A'X :: SA : AX$.

Alternately, $SA' : SA :: A'X : AX$,

$\therefore SA' - SA : SA :: A'X - AX : AX$,

or $AA' : SA :: XX' : AX$,

$\therefore AA' : XX' :: SA : AX$,

or $CA : CX :: SA : AX$. (1.)*

Again, $SA' : SA :: A'X : AX$.

$\therefore SA' + SA : SA :: A'X + AX : AX$,

or $SS' : SA :: AA' : AX$.

Alternately, $SS' : AA' :: SA : AX$,

or $CS : CA :: SA : AX$. (2.)

Hence, from (1) and (2),

$CA : CX :: CS : CA$,

$\therefore CA^2 = CX \cdot CS$.

Or CA is a mean proportional between CS and CX .

COR. Since the three lines CS , CA , CX , are proportional, therefore, by the definition of duplicate ratio, and *Euclid*, VI. 20 Cor.,

$CS : CX :: CS^2 : CA^2$. (3.)


PROP. III.

47. (If P be any point on the hyperbola, and S be the focus nearer to P ; then

$S'P - SP = AA'$.)

Since $SP : PM :: SA : AX$,

* N.B. The results (1), (2), (3), should be remembered, as they will be frequently referred to.



and $SA : AX :: AA' : XX'$, (*Prop. II.*)

$$\therefore SP : PM :: AA' : XY'.$$

So $S'P : PM' :: AA' : XX'$,

$$S'P - SP : PM - PM :: AA' : XX'.$$


but $PM' - PM = MM' = XX'$,

$$\therefore S'P - SP = AA'.$$

COR. By means of this property the hyperbola may be practically described, and the form of the curve determined.

Let a rigid bar $S'Q$ of any length have one end fastened at the focus S' , in such a manner that it is capable of turning freely round S' as a centre in the plane of the paper.

At the other end of the bar let a string be fastened of such a length that when stretched along the bar it shall just reach to within a distance equal to AA' from the end S' of the bar.



If the loose end of the string be now fastened to the focus S , and the rod being initially placed in the position $S'S$, be made to revolve round S' , while the string is kept constantly stretched by means of the point of a pencil at P , in contact with the bar; since $S'P$ and SP are always increasing by the same amount, viz. the length of the portion of the string that removes itself from the bar, between any two positions of P , the difference between $S'P$ and SP will be constantly the same, and the point P will trace out the hyperbola.

Another perfectly similar branch may be described in the same manner by making the bar revolve round S as centre.

In this case $SP - S'P$ will be equal to AA' .

The curve, therefore, consists of two similar branches, which recede indefinitely both from the line AA' , and also from the line BCB' drawn bisecting AA' at right angles. (See *fig. Prop. IV.*)

48. If BC be taken of such a length that

$$BC^2 = CS^2 - CA^2,$$

and CB' be made equal to CB , then AA' and BB' are called respectively the *Transverse* and *Conjugate Axes*.

The line BCB' does not meet the hyperbola, and the reason of its being introduced will be seen further on.

If the conjugate and transverse axes are equal, the hyperbola is said to be *rectangular* or *equilateral*.


The property of the hyperbola, which we have just investigated, viz. that the difference between SP and $S'P$ is constant, is sometimes taken as the definition of the curve. (See *Chapter II. Art. 25.*)

Also as in the ellipse, if SL be the semi-latus rectum, it may be proved that

$$SL \cdot AC = BC^2.$$

PROP. IV.


49. (The difference of the distances of any point from the foci of a hyperbola will be greater or less than AA' , according as the point is on the concave or convex side of the curve.)



(1.) Let Q be a point on the concave side of the hyperbola.

Join SQ , $S'Q$, and let $S'Q$ meet the curve in P ; join SP ; then

$$\begin{aligned} \text{since } S'Q &= S'P + PQ, \\ \text{and } SQ &< SP + PQ, \end{aligned}$$



$$\begin{aligned}\therefore S'Q - SQ &> S'P - SP. \\ \text{but } S'P - SP &= AA', \\ \therefore S'Q - SQ &> AA'.\end{aligned}$$

(2.) Let Q be a point on the convex side of the curve, nearer to S than S' ; join $S'Q$, SQ , and let SQ meet the curve in P ; join $S'P$; then

$$\begin{aligned}S'Q &< S'P + PQ, \\ \text{and } SQ &= SP + PQ, \\ \therefore S'Q - SQ &< S'P - SP, \\ \text{but } S'P - SP &= AA', \\ \therefore S'Q - SQ &< AA',\end{aligned}$$

so if Q be nearer to S' than S , we can show that

$$SQ - S'Q < AA';$$


COR. Conversely a point will be on the concave or convex side of the hyperbola, according as the difference of its distances from the foci is greater or less than AA' .

50. DEF. If a point P' be taken on the hyperbola near to P (see *fig. Prop. V.*), and PP' be joined, the line PP' produced, in the limiting position which it assumes when P' is made to approach indefinitely near to P , is called the *Tangent* to the hyperbola at the point P .

PROP. V.

(If the tangent to the hyperbola at any point P meet the directrix in the point Z , and if S be the focus corresponding to the directrix on which Z is situated, then SZ will be at right angles to SP .)

CONIC SECTIONS.



Let P' be a point on the curve near to P .

Draw the chord PP' , and produce it to meet the directrix in Z ; join SZ .

Draw PM , $P'M'$ at right angles to the directrix, and join SP , SP' .

Produce PS to meet the hyperbola in Q ; then since the triangles ZMP , $ZM'P'$ are similar,

$$\begin{aligned}\therefore ZP : ZP' &:: MP : M'P', \\ &:: SP : SP' .\end{aligned}$$

$\therefore SZ$ bisects the angle $P'SQ$. (*Euclid*, VI. A.)

Now, when P' is indefinitely near to P , and PP' becomes the tangent at the point P , the angle PSP' becomes indefinitely small, while the angle QSP' approaches two right angles; and therefore the angle ZSP' , being half of the angle $P'SQ$, becomes ultimately a right angle.

Hence, when PZ becomes the tangent at the point P , the angle ZSP is a right angle,

or SZ is perpendicular to SP .

COR. 1. Conversely, if SZ be drawn at right angles to SP , meeting the directrix in Z , and PZ be joined, PZ will be the tangent at P .


COR. 2. If PZ be produced to meet the other directrix in Z' , and $S'Z'$ be joined; then

$S'Z'$ is at right angles to $S'P$.

COR. 3. The tangents at the extremities of the latus rectum, or double ordinate through the focus, meet the axis in the point X .

PROP. VI.

(The tangent to the hyperbola at any point P makes equal angles with the focal distances SP and $S'P$.)



Let the tangent at P meet the directrices in Z and Z' .

Draw PMM' at right angles to the directrices meeting them in M and M' respectively; join SZ , $S'Z'$; then

$$SP : PM :: S'P : PM'.$$

And since the triangles $ZMP, Z'M'P$ are similar,

$$PM : PZ :: PM' : PZ',$$

$$\therefore SP : PZ :: S'P : PZ'. \quad (\text{Ex æquali.})$$

Now in the triangles $SPZ, S'PZ'$ because the sides about the angles $SPZ, S'PZ'$ are proportional, and the angles $PSZ, PS'Z'$ are equal, being right angles, and the angles $SZP, S'Z'P$ are each less than a right angle,

$\therefore$ the triangles $SPZ, S'PZ'$ are similar. (*Euclid*, VI. 7)

$$\therefore \text{the angle } SPZ = S'PZ'.$$

PROP. VII.

(The tangents at the extremities of a focal chord intersect in the directrix)

Let PSQ be a focal chord, and let the tangent at P meet the directrix in Z . Join SZ ; then

the angle ZSP is a right angle, (*Prop. V.*)

And $\therefore$ also the angle ZSQ is a right angle,

$\therefore ZQ$ is the tangent at Q . (*Prop. V. Cor. 1.*)

Or the tangents at the extremities of a focal chord intersect in the directrix.

PROP. VIII.

51. (If the tangent at P meet the transverse axis in T , and PN be the ordinate of the point P ; then

$$CT \cdot CN = CA^2.)$$

Draw PMM' at right angles to the directrices meeting them in M and M' . Join $SP, S'P$; then

since PT bisects the angle SPS' , (*Prop. VI.*)

$$\therefore S'T : ST :: S'P : SP, \quad (\text{Euclid, VI. 3.})$$

$$:: PM' : PM,$$

$$:: X'N : XN.$$

but the angle $S'PT =$ the angle $SP T$,
 $\therefore$ the angle $WPG =$ the angle SPG ,
 $\therefore PG$ bisects the angle SPW ,
 $\therefore S'G : SG :: S'P : SP$, (*Euclid*, VI. A.)
 $:: PM' : PM$,
 $:: X'N : XN$,
 $\therefore S'G + SG : S'G - SG :: X'N + XN : X'N - XN$;
 or $2CG : ES' :: 2CN : XX'$.
 Alternately, $2CG : 2CN :: SS' : XX$;
 or $CG : CN :: CS : CX$,
 $:: CS^2 : CA^2$ (*Prop. II. Cor.*)
 $\therefore CG - CN : CN :: CS^2 - CA^2 : CA^2$;
 or $NG : CN :: BC^2 : AC^2$.

PROP. X.

(If PN be the ordinate of any point I on the hyperbola,
 then

$$PN^2 : AN \cdot A'N :: BC^2 : AC^2.)$$

$$\text{For } NG : NC :: BC^2 : AC^2.$$

And rectangles of the same altitude are to one another as
 their bases, (*Euclid*, VI. 1.)

$$\therefore TN \cdot NG : TN \cdot NC :: BC^2 : AC^2,$$

$$\text{or } PN^2 : TN \cdot NC :: BC^2 : AC^2.$$

$$\text{But } TN \cdot CN = CN^2 - CT \cdot CN, \text{ (*Euclid*, II. 2.)}$$

$$= CN^2 - CA^2, \text{ (*Prop. VIII.*)}$$

$$= AN \cdot A'N, \text{ (*Euclid*, II. 6.)}$$

$$\therefore PN^2 : AN \cdot A'N :: BC^2 : AC^2.$$

PROP. XI.

If the normal at any point P of an hyperbola meet the
 transverse axis in G ; then

$$SG : SP :: CS : CA.$$

Produce $S'P$ to W ; then

since PG bisects the angle SPW , (*Prop. IX.*)

$$SG : S'G :: SP : S'P,$$

$$\therefore SG : S'G - SG :: SP : S'P - SP,$$

but $S'P - SP = AA'$, (*Prop. III.*)

and $S'G - SG = SS'$,

$$\therefore SG : SS' :: SP : AA',$$

$$\text{or } SG : SP :: SS' : AA',$$

$$\text{or } SG : SP :: CS : CA.$$

COR. Hence also,

$$S'G : S'P :: CS : CA.$$

PROP. XII.

53. If from the foci S and S' of an hyperbola SY and $S'Y'$ are drawn at right angles to the tangent at P , then Y and Y' are on the circumference of the circle described on AA' as diameter, and

$$SY \cdot S'Y' = BC^2.$$

Join SP , $S'P$, and produce SY to meet $S'P$ in W ; join CY ; then

since the angle $SPY =$ the angle WPY , (*Prop. VI.*)

and the angle $SY P =$ the angle WYP ,

and the side PY is common to the triangles SPY , WPY ,

$\therefore$ the triangle $SPY = WPY$ in all respects,

$$\therefore SP = PW, \text{ and } SY = WY,$$

$$\therefore S'P - SP = S'W,$$

but $S'P - SP = AA'$, (*Prop. III.*)


$$\therefore S'W = AA'.$$

Again, $\therefore SC = CS'$, and $SY = WY$,

$$\therefore SC : CS' :: SY : YW,$$

$\therefore CY$ is parallel to $S'W$,

$$\therefore CY : SW :: CS : SS',$$



$$\therefore CY = \frac{1}{2} S'W = CA;$$

$$\text{so } CY' = CA.$$

$\therefore Y$ and Y' are points on the circumference of the circle described upon AA' as diameter.

Next, let SY be produced to meet this circle in Z , and join ZY' ; then

since the angle ZYY' is a right angle

$\therefore ZY'$ passes through the centre C ,

$\therefore$ the angle $SCZ =$ the angle $S'CY'$,

$$\therefore SZ = S'Y',$$

$$\therefore SY \cdot S'Y' = SY \cdot SZ,$$

$$= SA \cdot SA', \text{ (Euclid, III. 36 Cor.)}$$

$$= CS^2 - CA^2, \text{ (Euclid, II. 6.)}$$

$$= BC^2.$$


COR. If CD be drawn parallel to the tangent at P meeting $S'P$ in E ; then

since $CEPY$ is a parallelogram,

$$\therefore PE = CY = AC.$$

PROP. XIII.

54. To draw a pair of tangents to an hyperbola from an external point O .



Of the foci S and S' , let S' be that which is nearer to O .

With centre S and radius equal to AA' describe a circle.

Join OS , OS' ; and let SO or SO produced meet the circle in the point I .

Now if O be a point inside the circle MIM' , it is evident that OS' is greater than OI ; and if O be outside the circle,

since $OS - OS' < AA'$ or SI , (*Prop. IV.*)

$\therefore OS - OS' < OS - OI$,

$\therefore OS' > OI$.

With centre O and radius OS' describe another circle cutting the former in the points M and M' , which it will always do since OS' is greater than OI .

Join SM, SM' , and produce them to meet the hyperbola in the points P and P' .

Join OP, OP' ; these will be the tangents required.

Join $S'P, S'P'$; then

$$\text{since } S'P - SP = AA' = SM,$$

$$\therefore S'P = PM.$$

And $\therefore S'P, PO = MP, PO$, each to each,

$$\text{and } OS' = OM,$$

$\therefore$ the angle $OPS' =$ the angle OPM ,

$\therefore OP$ is the tangent at P . (*Prop. VI.*)

So OP' is the tangent at P' .

The points of contact P and P' will be upon the same or opposite branches of the hyperbola according as SM and SM' require to be produced in the same or in opposite directions with respect to S , in order to intersect the hyperbola.

PROP. XIV.

If from a point O a pair of tangents, OP, OP' be drawn to an hyperbola, then the angles which OP and OP' subtend at either focus will be equal or supplementary according as the points of contact are in the same or opposite branches of the hyperbola.

Let the points P and P' be on opposite branches of the hyperbola.

Join $SP, S'P; SP', S'P'$.

Produce PS to M , making PM equal to PS' . Also from $P'S$ cut off a part $P'M'$ equal to $P'S'$.


Join $OM, OM'; OS, OS'$.

Then since $OP, PS' = OP, PM$, each to each,

and the angle $OPS' =$ the angle OPM , (*Prop. VI.*)

$$\therefore OS' = OM,$$

and the angle $OS'P =$ the angle OMP .



So $OS' = OM'$,
 and the angle $OS'P' =$ the angle $OM'P'$,
 $\therefore OM = OM'$.
 Again, $\because SM = S'P - SP = AA'$,
 and $SM' = SP' - S'P' = AA'$,
 $\therefore SM = SM'$.
 And $\because OS, SM = OS, SM'$, each to each,
 and $OM = OM'$,
 $\therefore$ the angle $OSM =$ the angle OSM' ,
 and the angle $OMS =$ the angle $OM'S$.
 But OSM is the supplement of OSP ,
 and $OM'S$ the supplement of $OM'P'$,
 $\therefore OSM'$ is the supplement of OSP ,
 and OMP the supplement of $OM'P'$.
 But $OMP = OS'P$,
 and $OM'P' = OS'P'$,
 $\therefore OS'P$ is the supplement of $OS'P'$.


Hence the angles which OP and OP' subtend either at S or S' are supplementary.

In a similar manner if P and P' are on the same branch of the hyperbola, the angles subtended either at S or S' may be shown to be equal.

PROP. XV.

55. If the tangent at any point P of an hyperbola meet the conjugate axis in the point t , and Pn be drawn at right angles to CB ; then

$$Cn \cdot Ct = BC^2.$$



Draw PN at right angles to CA ; then

$$Ct : CT :: PN : NT,$$

$$\therefore Ct : PN :: CT : NT,$$

$$\therefore Ct \cdot Cn : PN^2 :: CT \cdot CN : CN \cdot NT;$$

$$\text{or } Ct \cdot Cn : CT \cdot CN :: PN^2 : CN \cdot NT,$$

$$:: BC^2 : AC^2. \quad (\text{Prop. X.})$$

$$\text{But } CT \cdot CN = AC^2,$$

$$\therefore Ct \cdot Cn = BC^2.$$

56. The proofs that we have given up to this point of the properties of the hyperbola are closely analogous to the corresponding propositions in the ellipse. The remaining properties of the hyperbola are more conveniently investigated by means of its relation to certain lines, which we shall presently define, called *Asymptotes*, in the same manner as many of the properties of the ellipse were deduced from those of the auxiliary circle.

DEF. The hyperbola described (*see fig. Prop. XVI.*) with C as centre, and BB' as transverse axis, and AA' as conjugate axis, is called the *Conjugate Hyperbola*. Its foci, which will be on the line BCB' , will evidently be at the same distance from C as those of the original hyperbola, since

$$CS^2 = CA^2 + CB^2.$$

PROP. XVI.

If through any point R on either of the diagonals of the rectangle formed by drawing tangents to the hyperbola and its conjugate at the vertices, A, A', B, B' , two ordinates RPN, RDM , be drawn at right angles to AA' and BB' , and meeting either the hyperbola or its conjugate in the points P and D ; then

$$RN^2 \cup PN^2 = BC^2,$$

$$\text{and } RM^2 \cup DM^2 = AC^2.$$

Let R be a point on the diagonal $O'CO$; then

$$RN^2 : CN^2 :: AO^2 : AC^2,$$

$$:: BC^2 : AC^2;$$

$$\text{and } PN^2 : CN^2 - CA^2 :: BC^2 : AC^2; \text{ (Prop. X.)}$$

$$\therefore RN^2 - PN^2 : CA^2 :: BC^2 : AC^2.$$


$$\therefore RN^2 - PN^2 = BC^2.$$

$$\text{Again, } RM^2 : CM^2 :: AC^2 : BC^2,$$

$$\text{and } DM^2 : CM^2 - CB^2 :: AC^2 : BC^2; \text{ (Prop. X.)}$$

$$\therefore RM^2 - DM^2 : BC^2 :: AC^2 : BC^2$$

$$\therefore RM^2 - DM^2 = AC^2.$$



In exactly the same manner, if NR had been produced to meet the conjugate hyperbola in P , and MR had been produced to meet the original hyperbola in D , we should have had,

$$PN^2 - RN^2 = BC^2,$$

and $DM^2 - RM^2 = AC^2.$

COR. If RP be produced to meet the hyperbola in p , and the other asymptote in r ; then

$$RN^2 - PN^2 = RP \cdot Pr; \text{ (Euclid, II. 5.)}$$

$$\therefore RP \cdot Pr = BC^2.$$

Hence as RPN is further removed from A , and the line Pr consequently increases, since the rectangle contained by RP and Pr remains constant, RP must diminish, and by taking R sufficiently far from C , RP may be made less than any assignable magnitude. The line CR , therefore, continually approaches nearer and nearer to the hyperbola, though it never actually reaches it.

On account of this property, CR is called an *Asymptote* to the hyperbola.

So also if P be the point where NR produced meets the conjugate hyperbola, we shall have

$$RP \cdot Pr = BC^2;$$


and therefore CR is also the asymptote to the conjugate hyperbola.

In the same manner it may be shown that the other diameter oCo' of the rectangle OO' is an asymptote to both hyperbolas. ♣

PROP. XVII.

57. If E be the point where the asymptote meets the directrix, then

$$CE = AC.$$



For by similar triangles,

$$CE : CO :: CX : CA, \\ :: CA : CS. \quad (\text{Prop. II.})$$

$$\text{But } CO^2 = CA^2 + CB^2 = CS^2; \quad \checkmark$$

$$\therefore CO = CS;$$

$$\therefore CE = AC.$$

COR. If SE be joined, since

$$CE^2 = CA^2 = CS \cdot CX,$$

$\therefore$ the angle CES is a right angle. (*Euclid*, VI. 8, *Cor.*)

PROP. XVIII.

If from any point R in one of the asymptotes to an hyperbola ordinates RPN , $RD M$ be drawn to the hyperbola and its conjugate respectively, and PD be joined, PD will be parallel to the other asymptote.

$$\text{For } RN^2 : RM^2 :: BC^2 : AC^2;$$

$$\text{and } RN^2 - PN^2 : RM^2 - DM^2 :: BC^2 : AC^2, \quad (\text{Prop. XVI.})$$

$$\therefore PN^2 : DM^2 :: BC^2 : AC^2;$$

$$:: RN^2 : RM^2,$$

$$\therefore PN : DM :: RN : RM;$$

$$\therefore PD \text{ is parallel to } MN. \quad (\text{Euclid, VI. 2.})$$

$$\text{Also } CN : CM :: AC : BC,$$

$$\therefore MN \text{ is parallel to } AB;$$

$$\text{and } OA : Ao :: OB : Bo',$$

$$\therefore AB \text{ is parallel to } oo'.$$


Hence PD is parallel to oo' .

COR. So also if R and D be the points where NR and MR produced meet respectively the conjugate and the original hyperbola, PD will be still parallel to oo' .

PROP. XIX.

58. If through any two points Q and Q' of an hyperbola a line $RQ Q'R'$ be drawn in any direction meeting the asymptotes in R and R' ; then will

$$RQ = R'Q'.$$



Through Q and Q' draw the ordinates UQW , $U'Q'W'$; meeting the asymptotes in U , W , U' , W' ; then by similar triangles,

$$QR : QU :: Q'R : Q'U',$$

$$\text{and } QR' : QW :: Q'R' : Q'W';$$

$\therefore$ compounding

$$QR \cdot QR' : QU \cdot QW :: Q'R \cdot Q'R' : Q'U' \cdot Q'W'.$$

But $QU \cdot QW = BC^2 = Q'U' \cdot Q'W'$, (*Prop. XVI. Cor.*)

$$\therefore QR \cdot QR' = Q'R \cdot Q'R';$$

$$\begin{aligned}
&\text{but } QR \cdot QR' = QR \cdot QQ' + QR \cdot Q'R', \\
&\text{and } Q'R \cdot Q'R' = Q'R' \cdot QQ' + QR \cdot Q'R'; \\
&\therefore QR \cdot QQ' = Q'R' \cdot QQ', \\
&\therefore QR = Q'R'.
\end{aligned}$$

COR. 1. If $RQQ'R'$ move parallel to itself until the points Q and Q' coincide, the line RQR' will ultimately assume the position $LP l$, and will become a tangent to the hyperbola at P .

Hence, since RQ is always equal to $R'Q'$,

$$LP = Pl,$$

or the tangent $LP l$ is bisected at the point of contact P .

COR. 2. If CP be produced to meet RR' in V , then since

$$RV : VR' :: LP : Pl,$$

$$\therefore RV = VR';$$

$$\text{and } RQ = Q'R',$$

$$\therefore QV = Q'V.$$

Hence if a series of parallel chords be drawn in an hyperbola, their middle points will all be in the line drawn through the centre and the point where the tangent parallel to the chords meets the hyperbola.

DEF. A line PCP' drawn through the centre, and meeting the hyperbola in P and P' , is called a *Diameter*.


A diameter consequently bisects all chords drawn parallel to the tangents at its extremities.

PROP. XX.

59. If through any point Q of an hyperbola a line RQr be drawn in any direction meeting the asymptotes in R and r , and $LP l$ be the tangent drawn parallel to RQr ; then

$$RQ \cdot Qr = PL^2.$$

Through P and Q draw the ordinates EPe , UQW , meeting the asymptote in E , e , U , W ; then by similar triangles,



$$QR : QU :: PL : PE,$$

$$Qr : QW :: Pl : Pe,$$

$$\therefore QR \cdot Qr : QU \cdot QW :: PL \cdot Pl : PE \cdot Pe$$

$$\text{but } QU \cdot QW = BC^2 = PE \cdot Pe. \text{ (Prop. XVI. Cor.)}$$

$$\therefore QR \cdot Qr = PL \cdot Pl,$$

$$= PL^2. \text{ (Prop. XIX. Cor. 1.)}$$

COR. If Qq be produced to meet the conjugate hyperbola in Q', q' , we may show that

$$Q'R \cdot Q'r = PL^2,$$

and also, as in Proposition XIX., that


$$Q'R = q'r,$$

$$\therefore QQ' = qq'.$$

Hence if a line be drawn in any direction meeting both the hyperbolas, the portions intercepted between the hyperbola and its conjugate will be equal.

PROP. XXI.

60, If from any point P of an hyperbola, PH and PK be drawn parallel to the asymptotes, meeting them in H and K respectively; then $4 \cdot PH \cdot PK = CS^2$.



Draw the ordinate $RPNr$ meeting the asymptotes in R and r ; then by similar triangles,

$$PH : PR :: Co : Oo,$$

$$\text{and } PK : Pr :: CO : Oo,$$

$$\therefore PH \cdot PK : PR \cdot Pr :: CO^2 : Oo^2,$$

$$:: CS^2 : 4BC^2.$$

$$\text{But } PR \cdot Pr = BC^2, \quad \therefore$$

$$\therefore 4 \cdot PH \cdot PK = CS^2.$$

PROP. XXII.

If the tangent at any point P of an hyperbola meet the asymptotes in L and l ; then the area of the triangle LCl is equal to the rectangle contained by AC and BC .

Draw PH and PK parallel to the asymptotes meeting them in H and K ; then

$$\text{since } CL : CH :: Ll : Pl,$$

$$\text{and } Ll = 2 Pl, \quad (\text{Prop. XIX. Cor. 1.})$$

$$\therefore CL = 2 CH = 2 PK;$$

$$\text{so } Cl = 2 CK = 2 PH,$$

$$\therefore CL \cdot Cl = 4 PH \cdot PK = CS^2, \quad (\text{Prop. XXI.})$$

$$= CO \cdot Co,$$

$$\therefore CL : CO :: Co : Cl,$$

$\therefore$ the triangles LCl , OCo have the angle at C common and the sides about those angles reciprocally proportional;

$$\therefore \text{the triangle } LCl = \text{the triangle } OCo,$$

$$= AC \cdot AO,$$

$$= AC \cdot BC.$$

PROP. XXIII.

61. If from any point R in the asymptote of an hyperbola, two ordinates RPN and RDM be drawn to the hyperbola and its conjugate respectively, then the tangents at P and D will be parallel respectively to CD and CP .


Join PD , meeting CR in H ; then

$$\text{since } PD \text{ is parallel to } oo', \quad (\text{Prop. XVIII.})$$

the tangents at P and D will each meet CR produced in the same point L . (*Prop. XXII.*)

Produce LP and LD to meet the other asymptotes in l and l' ; then

$$\text{since } CL \cdot Cl = CS^2 = CL' \cdot Cl', \quad (\text{Prop. XXII.})$$



$$\therefore Cl = Cl',$$

$$\therefore lC : Cl' :: lP : PL,$$

$\therefore CP$ is parallel to the tangent at D .

$$\text{Also } l'D : DL :: l'C : Cl,$$

$\therefore CD$ is parallel to the tangent at P .

The lines CP and CD are called *Conjugate Diameters*, since each of these lines is parallel to the tangent at the extremity of the other.

PROP. XXIV.

If CP and CD be semi-conjugate diameters in the hyperbola; then

$$CP^2 \pm CD^2 = CA^2 \pm CB^2.$$


Draw the ordinates NPR , MDR meeting the asymptote in the point R (*Prop. XXIII.*); then

H

$$\begin{aligned}
 CR^2 - CP^2 &= NR^2 - NP^2, \\
 &= BC^2, \quad (\text{Prop. XVI.}) \\
 \therefore CR^2 &= CP^2 + BC^2; \\
 \text{so } CR^2 &= CD^2 + AC^2, \\
 \therefore CP^2 + BC^2 &= CD^2 + AC^2; \\
 \text{or } CP^2 \hookrightarrow CD^2 &= AC^2 \hookrightarrow BC^2.
 \end{aligned}$$

PROP. XXV.

62. The area of any parallelogram formed by drawing tangents to the hyperbola and its conjugate at the extremities P, P', D, D' of a pair of conjugate diameters is equal to the rectangle contained by the axes.



Let $LL'L'l'$ be the parallelogram formed by drawing tangents at the extremities P, P', D, D' , of any pair of conjugate diameters. The points L, L', l, l' , will (*Prop. XXIII.*) be on the asymptotes.

Now the parallelogram $LL' = 4$ parallelogram CL ,
 $= 4$ triangle LCl ,
 $= 4AC \cdot BC$, (*Prop. XXII.*)
 $= AA' \cdot BB'$.

COR. If PF be drawn perpendicular to CD , then
 $PF \cdot CD = AC \cdot BC$.

Also, if the normal PG meet the transverse axis in G , as in the ellipse

$$PF \cdot PG = BC^2.$$

63. DEF. The line QV drawn from any point Q of the hyperbola parallel to the tangent at any point P , and meeting CP produced in V , is called an *Ordinate* to the diameter CP .

PROP. XXVI.

If QV be an ordinate to the diameter $P'CP$, and CD be conjugate to CP ; then .

$$QV^2 : PV \cdot P'V :: CD^2 : CP^2.$$

Produce VQ to meet the asymptotes in R and r ; and let the tangent at P meet the asymptotes in L and l ; then

$$RV^2 : PL^2 :: CV^2 : CP^2,$$

$$\therefore RV^2 - PL^2 : PL^2 :: CV^2 - CP^2 : CP^2.$$

But $RQ \cdot Qr = PL^2$, (*Prop. XX.*)

$$\therefore RV^2 - QV^2 = PL^2,$$

$$\text{or } RV^2 - PL^2 = QV^2.$$

And $CV^2 - CP^2 = PV \cdot P'V$, (*Euclid, II. 6*)

$$\therefore QV^2 : PL^2 :: PV \cdot P'V : CP^2.$$

Alternately, $QV^2 : PV \cdot P'V :: PL^2 : CP^2$.

But since PD is a parallelogram, (*Prop. XXIII.*)

$$\therefore PL = CD.$$

Hence $QV^2 : PV \cdot P'V :: CD^2 : CP^2$.

COR. If VQ be produced to meet the conjugate hyperbola in Q' , then

since $Q'R \cdot Q'r = PL^2$, (*Prop. XX. Cor.*)

$$\therefore Q'V^2 - RV^2 = PL^2.$$


Hence $Q'V^2 : CV^2 + CP^2 :: CD^2 : CP^2$

PROP. XXVII.

64. If QV be an ordinate to the diameter PV , and the tangent at Q meet CP in the point T ; then

$$CV \cdot CT = CP^2.$$

Draw the tangent LPl meeting the asymptotes in the points L, l ; also let the tangent at Q meet the asymptotes in R, r .



Draw RK, rk parallel to QV meeting CP in K, k .

Now since the triangles RCr, LCl are equal, (*Prop. XXII.*) and have the angle at C common,

$$\therefore CR : CL :: Cl : Cr. \text{ (Euclid, VI. 15.)}$$

$$\text{But } CR : CL :: CK : CP,$$

$$\begin{aligned}
 &\text{and } Cl : Cr :: CP : Ck, \\
 &\therefore CK : CP :: CP : Ck, \\
 &\therefore CK \cdot Ck = CP^2.
 \end{aligned}$$

Again, produce RK and QV to meet the asymptote Cl in R' and q ; then

Since Rr is bisected in Q , (*Prop. XIX. Cor. 1*)

$\therefore R'r$ is bisected in q ,
and $RK = R'K$, (*Prop. XIX. Cor. 2*)

$$\begin{aligned}
 &\therefore Kq \text{ is parallel to } Rr, \\
 &\therefore CT : CK :: Cr : Cq, \\
 &\qquad\qquad\qquad :: Ck : CV, \\
 &\therefore CV \cdot CT = CK \cdot Ck \\
 &\qquad\qquad\qquad = CP^2.
 \end{aligned}$$

COR. 1. Conversely, if QV be an ordinate to PV ,

and $CV \cdot CT = CP^2$,

then QT is the tangent at Q .

COR. 2. Hence also, if RR' meet the curve in U and U' ,

and kU, kU' be drawn,

since $CK \cdot Ck = CP^2$,

$\therefore kU$ and kU' are tangents to the hyperbola at U and U' .


PROP. XXVIII.

65. If two chords of a hyperbola intersect one another, the rectangles contained by their segments are proportional to the squares of the diameters parallel to them.

Let QOq be any chord drawn through the point O , and let CD be drawn parallel to it, meeting the conjugate hyperbola in D .

Produce Qq to meet the asymptotes in R and r ; and draw the diameter CPV , bisecting both Qq and Rr in V . (*Prop. XIX. Cor. 2.*)

Also draw the tangent LPl parallel to Qq , meeting the asymptotes in L and l .



Now since Qq is divided equally in V and unequally in O ,

$$\therefore QO \cdot Oq = QV^2 - OV^2; \text{ (Euclid, II. 5)}$$

$$\text{so also } RO \cdot Or = RV^2 - OV^2, \text{ (Euclid, II. 5)}$$

$$\begin{aligned} \therefore RO \cdot Or - QO \cdot Oq &= RV^2 - QV^2, \\ &= RQ \cdot Qr \text{ (Euclid, II. 5)} \\ &= PL^2, \text{ (Prop. XX.)} \end{aligned}$$

$$\therefore QO \cdot Oq = RO \cdot Or - PL^2.$$

Again, through O and P draw EOe , UPW , at right angles to the axis meeting the asymptotes in E , e , U , W ; then

$$RO : OE :: PL : PU,$$

$$\text{and } rO : Oe :: Pl : PW,$$

$$\therefore RO \cdot rO : OE \cdot Oe :: PL^2 : PU \cdot PW;$$

but $P\dot{U} \cdot PW = BC^2$, (*Prop.* XVI.)

and $PL^2 = CD^2$, (*Prop.* XXIII.)

$$\therefore RO \cdot rO : OE \cdot Oe :: CD^2 : BC^2,$$

$$\text{or } R\dot{O} \cdot rO : CD^2 :: OE \cdot Oe : BC^2,$$

$$\therefore RO \cdot rO - PL^2 : CD^2 :: OE \cdot Oe - BC^2 : BC^2,$$

$$\text{or } QO \cdot Oq : CD^2 :: OE \cdot Oe - BC^2 : BC^2.$$

In the same manner if through O another chord $Q'Oq'$ be drawn, and CD' be drawn parallel to it, meeting the conjugate hyperbola in D' , we shall have

$$Q'\dot{O} \cdot Oq' : CD'^2 :: OE \cdot Oe - BC^2 : BC^2.$$

$$\text{Hence } QO \cdot Oq : Q'O \cdot Oq' :: CD^2 : CD'^2.$$

COR. The same result may be shown to be true when the point O is outside the hyperbola. Moreover, it is not necessary that the chords should be drawn meeting *one* branch only of the hyperbola or the *same* branch. The proportion still holds good when one or both of the chords meet both branches of the hyperbola, or when the chords are drawn in different branches.

66. DEF. If with a point O on the normal at P as centre, and OP as radius, a circle be described touching the hyperbola at P , and cutting it in Q ; then when the point Q is made to approach indefinitely near to P , the circle is called the *Circle of Curvature* at the point P .

PROP. XXIX.

If PH be the chord of the circle of curvature at the point P of a hyperbola, which passes through the centre; then


$$PH \cdot CP = 2CD^2.$$

Let PT be the tangent, and PG the normal at the point P .

With centre O , and radius OP , describe a circle cutting the hyperbola in the point Q .

Draw RQW parallel to CP , meeting the circle in W , and TP produced in R .

Also, draw QV parallel to PR , meeting the diameter PP' in V ; then since RP touches the circle at P ,



$$\therefore RQ \cdot RW = PR^2, \text{ (Euclid, III. 36)}$$

$$\text{or } PV \cdot RW = QV^2.$$

But $QV^2 : PV \cdot P'V :: CD^2 : CP^2$, (Prop. XXVI.)

$$\therefore PV \cdot RW : PV \cdot P'V :: CD^2 : CP^2,$$

$$\text{or } RW : P'V :: CD^2 : CP^2,$$

Now, when the circle becomes the circle of curvature at P , the points R and Q move up to, and coincide with P , and the lines RW and PH become equal, while

$$P'V \text{ becomes equal to } PP', \text{ or } 2CP.$$

$$\text{Hence, } PH : 2CP :: CD^2 : CP^2,$$

$$\therefore PH \cdot CP : 2CP^2 :: 2CD^2 : 2CP^2,$$

$$\therefore PH \cdot CP = 2CD^2.$$

PROP. XXX.

If PU be the diameter of the circle of curvature at the point P of the hyperbola, and PF be drawn at right angles to CD ; then

$$PU \cdot PF = 2CD^2.$$

Since the triangle PHU is similar to the triangle PFC ,

$$\therefore PU : PH :: CP : PF,$$

$$\therefore PU \cdot PF = PH \cdot CP, \\ = 2 CD^2. \quad (Prop. XXIX.)$$

PROP. XXXI.

If PI be the chord of the circle of curvature through the focus of the hyperbola; then

$$PI \cdot AC = 2 CD^2.$$

Let $S'P$ meet CD in E ; then since the triangles PIU and PEF are similar,

$$\therefore PI : PU :: PF : PE$$

$$\text{But } PE = AC, \quad (Prop. XII. Cor.)$$

$$\therefore PI : PU :: PF : AC,$$

$$\therefore PI \cdot AC = PU \cdot PF, \\ = 2 CD^2. \quad (Prop. XXX.)$$

The point where the circle of curvature intersects the hyperbola may be determined as in the case of the ellipse.

PROP. XXXII.


67. If P be any point on the hyperbola, and CD be conjugate to CP ; then

$$SP \cdot S'P = CD^2.$$

Draw PII' parallel to the asymptote CE meeting the directrices in I and I' , and CB' in U .

Let the ordinates NP , MD meet the asymptote in R , and draw PW perpendicular to the directrix; then by similar triangles,

$$PI : PW :: CE : CX, \\ :: CA : CX. \quad (Prop. XVII.)$$



But $SP : PW :: SA : AX,$
 $:: CA : CX.$

$\therefore SP = PI;$

so $S'P = PI',$

$\therefore SP \cdot S'P = PI \cdot PI',$
 $= UP^2 - UI^2,$
 $= CR^2 - CE^2,$
 $= CR^2 - CA^2. (Prop. XVII.)$

But $CR^2 - CD^2 = RM^2 - DM^2,$
 $= CA^2, (Prop. XVI.)$

$\therefore CR^2 - CA^2 = CD^2.$

Hence $SP \cdot S'P = CD^2.$

PROBLEMS ON THE HYPERBOLA.

1. THE locus of the centre of a circle touching two given circles is a hyperbola or ellipse.

2. If on the portion of any tangent intercepted between the tangents at the vertices a circle be described, it will pass through the foci.

3. In a hyperbola the tangents at the vertices will meet the asymptotes in the circumference of the circle described on SS' as diameter.

4. If from a point P in a hyperbola PII' be drawn parallel to the transverse axis meeting the asymptotes in I and I' ; then $PI \cdot PI' = AC^2$.

5. If a circle be inscribed in the triangle SPS' , the locus of its centre is the tangent at the vertex.

6. If PN be the ordinate of the point P , and NQ a tangent to the circle described on the transverse axis as diameter, and PM be drawn parallel to QC meeting the axis in M , then $MN = BC$.

7. If PN be the ordinate of a point P , and NQ be drawn parallel to AP to meet CP in Q , then AQ is parallel to the tangent at P .

8. If a hyperbola and an ellipse have the same foci, they cut one another at right angles.

9. If the tangent at P intersect the tangents at the vertices in R, r , and the tangent at P' intersect them in R', r' , then $AR \cdot Ar = AR' \cdot Ar'$.

10. If any two tangents be drawn to a hyperbola, the lines joining the points where they intersect the asymptotes will be parallel.

11. The perpendicular drawn from the focus to the asymptotes of a hyperbola is equal to the semi-conjugate axis.

12. If the asymptotes meet the tangent at the vertex in O , and the directrix in E ; then AE is parallel to SO .

13. In a rectangular hyperbola conjugate diameters are equal to one another.

14. In a rectangular hyperbola the normal PG is equal to CP .

15. The lines drawn from any point in a rectangular hyperbola to the extremities of a diameter make equal angles with the asymptotes.

16. Prove that the asymptotes to a hyperbola bisect the lines joining the extremities of conjugate diameters.

17. A line drawn through one of the vertices of a hyperbola and terminated by two lines drawn through the other vertex parallel to the asymptotes will be bisected at the other point where it cuts the hyperbola.

18. P is any point on a hyperbola, and P' a point on the conjugate hyperbola. If CP and CP' be conjugate, prove that

$$S'P' - SP = AC - BC,$$

S and S' being the interior foci.

19. If CP and CD be conjugate, and through C a line be drawn parallel to either focal distance of P , the perpendicular from D upon this line is equal to BC .

20. Given a pair of conjugate diameters, find the principal axes.

21. If Q be a point on the conjugate axis of a rectangular hyperbola, and QP be drawn parallel to the transverse axis meeting the curve in P ; then

$$PQ = AQ.$$

22. In a rectangular hyperbola the focal chords drawn parallel to conjugate diameters are equal.

23. If in an equilateral hyperbola CY be drawn at right angles to the tangent at P , and AY be joined, the triangles PCA , CAY are similar.

24. The radius of the circle which touches a hyperbola and its asymptotes, is equal to that part of the latus rectum produced which is intercepted between the curve and the asymptotes.

25. If QQ' be any chord of a hyperbola, and CP the diameter corresponding to it, and QH , PK , $Q'H'$ be drawn parallel to one asymptote meeting the other in H , K and H' , then $CH \cdot CH' = CK^2$.

26. If the chord $RPP'R'$ intersect the hyperbola in the points P , P' , and the asymptotes in R , R' ; and PK be drawn parallel to CR' , and $P'K'$ to CR ; then $RK = P'K'$, and $R'K' = PK$.

27. If AA' be any diameter of a circle, and PNQ an ordinate to it, then the locus of the intersections of AP , $A'Q$ is a rectangular hyperbola.

28. If two concentric rectangular hyperbolas be described, the axes of one being the asymptotes of the other, they will intersect at right angles.

29. If any chord AP through the vertex be divided in Q , so that $AQ : QP :: AC^2 : BC^2$, and QN be drawn to the foot of the ordinate PN , prove that a straight line drawn at right angles to QN from Q cuts the transverse axis in the same ratio.

30. Prove that the curve which trisects the arcs of all segments of a circle described upon a given base is a hyperbola.

31. If SVs , TVt be two tangents cutting one asymptote in S , T , and the other in s , t , prove that

$$VS : Vs : Vt : VT.$$

32. If from the exterior focus of a hyperbola a circle be described with radius equal to BC , and tangents be drawn to it from any point in the hyperbola, the line joining the points of contact will touch the circle described on the transverse axis as diameter.

33. Circles are drawn touching the straight line AB in a fixed point C ; and from the fixed points A, B tangents are drawn to these circles. The locus of their intersection is an ellipse or hyperbola. Distinguish between the two cases.

34. PP' is a double ordinate in an ellipse. $AP, A'P'$ are produced to meet in Q . Prove that the locus of Q is a hyperbola with the same axes as the ellipse.

35. *on a rectangular hyperbola.* If the tangent at P intersect the asymptotes in L and l , and PG be the normal at P , then the angle LGl is a right angle.

36. If an ellipse, a parabola, and a hyperbola, have a common tangent, and the same curvature at the vertex, the ellipse will be entirely within the parabola, and the parabola entirely within the hyperbola.

37. The chord $RPP'R'$ of a hyperbola intersects the asymptotes in R and R' . From the point R a tangent RQ is drawn meeting the hyperbola in Q . If $PH, QK, P'H'$ be drawn parallel to one asymptote meeting the other in the points H, K, H' ; then $PH + P'H' = 2 QK$.

38. If through P, P' on a hyperbola lines be drawn parallel to the asymptotes forming a parallelogram, of which PP is one diagonal; the other diagonal will pass through the centre.

39. If P be the middle point of a line EF which moves so as to cut off a constant area from the corner of a rectangle, its locus is an equilateral hyperbola.

40. PM, PN are drawn parallel to the asymptotes CN, CM , and an ellipse is constructed having CN, CM for semi-conjugate diameters. If CP cut the ellipse in Q , the tangents at Q and P to the ellipse and hyperbola are parallel.

41. If a circle be described through any point P of a given hyperbola and the extremities A, A' of the transverse axis, and NP be produced to meet the circle in Q ; prove that Q traces out a hyperbola whose conjugate axis is a third proportional to the conjugate and transverse axes of the original hyperbola.

42. If lines be drawn from any point of a rectangular hyperbola to the extremities of a diameter, the difference between the angles which they make with the diameter will be equal to the angle which this diameter makes with its conjugate.

43. If between a rectangular hyperbola and its asymptotes any number of concentric elliptic quadrants be inscribed, the rectangle contained by their axes will be constant.

44. In the rectangular hyperbola if CP be produced to Q so that $PQ = CP$, and QO be drawn at right angles to CQ to intersect the normal in O , O is the centre of curvature at P .

45. With two conjugate diameters of an ellipse as asymptotes a pair of conjugate hyperbolas are constructed; prove that if one hyperbola touch the ellipse the other will do so likewise; prove also that the diameters drawn through the points of contact are conjugate to each other.

46. If a pair of conjugate diameters of an ellipse when produced be asymptotes to a hyperbola, the points of the hyperbola at which a tangent to the hyperbola will also be a tangent to the ellipse, lie in an ellipse similar to the given one.

47. In the rectangular hyperbola the radius of curvature at P is to the radius of curvature at P' in the triplicate ratio of CP to CP' .

48. OP , OQ are tangents to an ellipse at P and Q , and asymptotes to a hyperbola. Show that a pair of their common chords is parallel to PQ . One of these chords being RS , prove that if PR touches the hyperbola at P , QS touches it at S ; also if PS , QR meet in U , OU bisects PQ .

49. The base of the triangle ABC remains fixed, while the vertex C moves in an equilateral hyperbola passing through A and B . If P , Q be the points in which AC , BC meet the circle described on AB as diameter, the intersection of AQ , BP is on the other branch of the hyperbola.

CHAPTER IV.

THE SECTIONS OF THE CONE.

68. DEF. If two indefinite straight lines IOI' , DOD' , intersect one another at a point O , and one of them IOI' remain fixed while the other DOD' revolves round it in such a manner that its inclination to IOI' is the same in all positions, the surface generated by DOD' will be a *Right Cone*.


The line IOI' is called the *Axis*, and the point O the *Vertex* of the *Cone*.

It now remains for us to show (*see Introduction*) that the curve formed by the intersection of this surface with a plane is *in general* one of the three curves whose properties we have been investigating, and to consider under what circumstances it will be the Parabola, Ellipse, or Hyperbola.

If the cutting plane pass through the vertex of the cone as KOK' , and intersect the cone again *at all*, it will *in general* cut it in two straight lines as OK , OK' , which will represent two positions of the generating line.

The inclination of these lines to each other will depend upon the inclination of the cutting plane to the axis of the cone, and will be greatest when this plane passes through the axis, in which case it will be double the constant angle between the axis and the generating line.

If the cutting plane pass through a generating line dod' and be perpendicular to the plane containing this line and the axis, it will simply touch the cone along this line.



Should the cutting plane not pass through the vertex, and be at right angles to the axis of the cone, the section will evidently be a circle.

In any other case the section will, as we proceed to show, be a Parabola, Ellipse, or Hyperbola.

Whatever be the position of the cutting plane with respect to the cone, we can always suppose a plane drawn through the axis of the cone at right angles to it; and it will be convenient to have this latter plane represented by the plane of the paper as DOd . The cutting plane will therefore always be taken at right angles to the plane DOd of the paper.

PROP. I.

69. The curve formed by the intersection of the surface of a right cone with a plane (which neither passes through its vertex nor is at right angles to its axis) will be a *Parabola*, *Ellipse*, or *Hyperbola*, according as the inclination of the cutting plane to the axis of the cone is *equal to*, *greater*, or *less than* the constant angle which the generating line forms with the axis.

Let the plane of the paper represent the plane drawn through the axis IOI' of the cone at right angles to the cutting plane; and let it intersect the surface of the cone in the two generating lines OD , Od .

Let the cutting plane intersect the surface of the cone in the curve PA , and the plane of the paper in the line ANH .

The curve will evidently be symmetrical with respect to this line.

On AH take any point N , and through N draw a plane perpendicular to the axis meeting the surface of the cone in the circle $RP r$, and the cutting plane in the line PN , which will be at right angles to the plane of the paper and to AN .

Let a sphere be inscribed in the cone touching the cone in the circle EQe and the cutting plane in the point S , and let the plane EQe intersect the cutting plane in the line XM , which will be at right angles to the plane of the paper, and therefore parallel to PN .

Draw PM perpendicular to XM , and join PS , PO , and let PO meet the circle EQe in the point Q .*


Then since PS and PQ are both tangents to the sphere,

$$\therefore PS = PQ.$$

$$\text{But } PQ = RE,$$

$$\therefore PS = RE.$$

* N.B. In the figure, to avoid confusion, that part of the section which lies above the plane of the paper is alone represented.



But $RE : XN :: AE : AX$, (*Euclid*, VI. 2.)

and $AE = AS$,

$\therefore RE : XN :: AS : AX$,

$\therefore SP : PM :: AS : AX$,

$\therefore$ the curve PA is either a *Parabola*, *Ellipse*, or *Hyperbola*, whose focus is S and directrix XM .

Again, let AH meet the axis OI in F .

Then the angle AFO will be the inclination of the cutting plane to the axis.

(1.) Let the angle $AFO =$ the angle $F Od$;


then AH is parallel to Od ,

$\therefore$ the angle $AXE =$ the angle $OeE =$ the angle AEX ,

$\therefore AE = AX$,

$\therefore AS = AX$,

$\therefore$ the curve AP will be a *Parabola*.



- (2.) Let the angle AFO be $> FOd$; then
 the complement FXE is $<$ the complement $O Ee$,
 $\therefore$ the angle AXE is $<$ the angle AEX ,
 $\therefore AE$ is $< AX$,
 or AS is $< AX$,
 $\therefore$ the curve AP is an *Ellipse*.

Since the angles HFO , FOd are together less than HFO , OFA , i.e. than two right angles, the lines AH and Oe may be produced to meet in A' .

If another sphere be described touching the cone in the circle $E'Q'e'$ and the cutting plane in the point S' ; and the line $X'M'$ denote the intersection of the plane $E'Q'e'$ with the cutting plane, and PM' be drawn at right angles to this line, it can easily be shown that

$$S'P : PM' :: S'A' : A'X'.$$

Hence S' and $X'M'$ represent respectively the other focus and directrix of the ellipse.

Also if BC be the semi-axis minor, and through the centre C a line UCU' be drawn parallel to Ee meeting OD , Od in U and U' , then it is evident that

$$BC^2 = CU \cdot CU'.$$

- (3.) Let the angle AFO be $< FOd$; then
 the angle AXE is $>$ the angle AEX ,
 $\therefore AE$ is $> AX$,
 $\therefore AS$ is $> AX$,
 $\therefore$ the curve PA is an *Hyperbola*.

Since the angles AFO , FOd' are less than the two FOd , FOd' , i.e. than two right angles, the lines FA and dO may be produced to meet in A' .

In this case the cutting plane will intersect the other half of the cone, and if any point P' be taken on this part of the curve, and $P'M$ be drawn at right angles to XM , it can be shown as before that

$$SP' : P'M :: SA : AX.$$

The intersection of the cutting plane therefore with this portion of the cone will be the other branch of the hyperbola.

Also if another sphere be described touching the upper portion of the cone in $E'Q'e'$, and the cutting plane in S' , and the line $X'M'$ denote the intersection of the plane $E'Q'e'$ with the cutting plane, and $P'M'$ be drawn at right angles to this line, it can be easily shown that

$$S'P' : P'M' :: S'A' : A'X'.$$

Hence S' and $X'M'$ will represent respectively the other focus and directrix of the hyperbola.

COR. 1. In this last case, *i.e.* when the section is an hyperbola, if a plane OKL be drawn through the vertex of the cone parallel to the cutting plane, meeting the plane of the paper in the straight line OL , and the surface of the cone in the generating line OK ; then

$$\begin{aligned} OL : OK &:: OL : OR, \\ &:: AN : AR, \\ &:: AX : AE, \text{ (Euclid, VI. 2)} \\ &:: AX : AS, \\ &:: CA : CS, \text{ (Chap. III. Prop. II.)} \end{aligned}$$

where C is the middle point of AA' , and therefore the centre of the hyperbola.

$\therefore KOL$ is half the angle between the asymptotes. (*Chap. III. Prop. XVI.*)

Again, if BC be the conjugate semi-axis, and $CU'U$ be drawn parallel to Rr meeting OD' , Od' in U and U' , then

$$\begin{aligned} \text{since } CU : AC &:: RL : OL, \\ \text{and } CU' : A'C &:: rL : OL, \\ \therefore CU \cdot CU' : AC^2 &:: RL \cdot rL : OL^2, \\ &:: KL^2 : OL^2; \\ \text{but } BC^2 : AC^2 &:: KL^2 : OL^2, \\ \therefore BC^2 &= CU \cdot CU'. \end{aligned}$$

COR. 2. If the cutting plane is parallel to the axis, OL and OI coincide.

PROBLEMS ON THE SECTIONS OF THE CONE.

1. THE foci of all parabolic sections which can be cut from a given right cone, lie upon the surface of another cone.

2. The foci of all elliptical sections of a given right cone, in which the ratio of CA to CS is the same, will lie on two other cones.

3. The extremities of the minor axes of the elliptical sections of a right cone made by parallel planes, lie on two generating lines.

4. The latus rectum of a parabola cut from a given cone varies as the distance between the vertices of the cone and the parabola.

5. Under what conditions is it possible to cut an equilateral hyperbola from a given right cone?

6. Two cones whose vertical angles are supplementary are joined as in *Art. 69, Cor. 2*. Prove that the latera recta of the curves of section of the cones, whose axes are respectively OI and Oi , made by planes parallel or perpendicular to the plane of the axes, are in the duplicate ratio of $Oi : OI$.

ADDITIONAL PROBLEMS.

1. SHOW that the part of the directrix of a parabola, intercepted between the perpendiculars on it from the extremities of any focal chord, subtends a right angle at the focus.

2. The locus of the foci of all parabolas touching the three sides of a triangle is a circle. Prove this, and give a geometrical construction for finding the centre.

3. A system of parabolas which always touch two given straight lines have their axes parallel; show that the locus of the foci is a straight line.

4. Prove that the locus ~~at~~^{of} the foot of the perpendicular from the focus of a parabola on the normal is a parabola.

5. If S be the focus of a parabola which touches the sides AB , AC of the triangle ABC at the points B , C , and O the centre of the circle described about the triangle; prove that the angle OSA is a right angle.

6. From the focus of a parabola a straight line is drawn parallel to the tangent at any point P , meeting the diameter through P in V' ; show that the tangent drawn from P to any circle passing through V' is equal to one-half of the ordinate QV , V being the second point in which the circle cuts the diameter through P .

7. PSp is a focal chord, and upon PS and pS as diameters, circles are described; prove that the length of either of their common tangents is a mean proportional between AS and Pp .

8. If AQ be a chord of a parabola through the vertex A , and QR be drawn perpendicular to AQ to meet the axis in R ; prove that AR will be equal to the chord through the focus parallel to AQ .

9. The locus of the vertices of all parabolas, which have a common focus and a common tangent, is a circle.

10. Two parabolas have a common axis and vertex, and their concavities turned in opposite directions; the latus rectum of one is eight times that of the other; prove that the portion of a tangent to the former, intercepted between the common tangent and axis, is bisected by the latter.

11. B is a point on a radius OA of a circle, whose centre is O . On OA produced a point C is taken, such that $OB \cdot OC = OA^2$. If P be any point on the circumference of this circle, R the middle point of BP , and Q the point of intersection of AR , CP ; prove that the locus of Q is a circle.

12. If from the middle point of a focal chord of a parabola two straight lines be drawn, one perpendicular to the chord meeting the axis in G , and the other perpendicular to the axis meeting it in N ; show that NG is constant.

13. A circle is drawn touching the axis of a parabola, the focal distance of a point P , and the diameter through P . Show that the locus of its centre is a parabola with vertex S , and latus rectum equal to AS .

14. If from the point of intersection of the directrix and axis of a parabola a chord XPQ be drawn, cutting the parabola in P , Q ; show that the rectangle contained by the ordinates of PQ is equal to the square of one-half the latus rectum.

15. Find the locus of the centre of a circle which touches a given circle and a given straight line.

16. Given one point of contact of a parabola with three

tangents given in position, find the two other points of contact.

17. The triangle ABC circumscribes a parabola whose focus is S . Through A, B, C , lines are drawn perpendicular respectively to SA, SB, SC . Show that these lines pass through one point.

18. From the focus a line is drawn parallel to the tangent at P , meeting the parabola at Q . QN is an ordinate, and the tangents at P and Q meet the axis in T and T' . Prove that $SN^2 = 4AT \cdot AT'$, and that if the diameter at P meet SQ in E , the locus of E is a parabola, whose latus rectum is half that of the given parabola.

19. P is any point in a parabola; through S a line is drawn at right angles to the axis, meeting the chord AP or AP produced in R . Prove that $SK \cdot SR = 2SY^2$, where SY is the perpendicular on the tangent, and SK on the normal.

20. From the focus S of a parabola SK is drawn, making a given angle with the tangent at P . Show that the locus of K is that tangent to the parabola which makes with the axis an angle equal to the given angle.

21. PSQ is a focal chord of a parabola, AP' a parallel chord meeting the latus rectum in Q' ; prove that $AP' \cdot AQ' = SP \cdot SQ$.

22. The circle of curvature at any point of a parabola whose abscissa is AN cuts the axis in U and U' . Prove that $AU \cdot AU' = 3AN^2$.

23. AB is a diameter of a circle. From any point Q in the circumference a tangent QP is drawn, and from P a perpendicular PN is let fall upon AB . Show that if P be always taken so that QP is equal to AN , the locus of P will be a parabola.

24. If a tangent be drawn from any point of a parabola to the circle of curvature at the vertex, the length of the tangent

will be equal to the abscissa of the point measured along the axis.

25. To two parabolas which have a common focus and axis two tangents are drawn at right angles; the locus of their intersection is a straight line parallel to the directrices.

26. If any three tangents be drawn to a parabola, the circle described about the triangle so formed will pass through the focus, and the perpendiculars from the angles on the opposite sides intersect in the directrix.

27. A parabola touches one side of a triangle in its middle point, and the other two sides produced. Prove that the perpendicular drawn from the angles of the triangle upon any tangent to the parabola are in harmonical progression.

28. Two equal parabolas have the same axis and vertex, but are turned in opposite directions; chords of one parabola are tangents to the other. Show that the locus of the middle point of the chords is a parabola whose latus rectum is one-third of that of the given parabola.

29. Two equal parabolas have the same focus, and their axes are at right angles to each other, and a normal to one of them is perpendicular to a normal of the other; prove that the locus of the intersection of such lines is a parabola.

30. Show that in every ellipse there are two equal conjugate diameters, coinciding in direction with the diagonals of the rectangle, which touches the ellipse at the extremities of the axes.

31. If a circle be described through the two foci of an ellipse, cutting the ellipse, show that the angle between the tangents to this circle, and to the ellipse at either point of intersection, is equal to the inclination of the normal to the ellipse to the axis minor.

32. The points in which the tangents at the extremities of the transverse axis of an ellipse are cut by the tangent at any

point of the curve, are joined one with each focus; prove that the point of intersection of the joining lines lies in the normal at the point.

33. The external angle between any two tangents to an ellipse is equal to the semi-sum of the angles which the chord joining the points of contact subtends at the foci.

34. The tangent to an ellipse at any point P is cut by any two conjugate diameters in T, t , and the points T, t are joined with the foci S, S' respectively; prove that the triangles $SPT, S'Pt$ are similar to each other.

35. P is any point on a fixed circle, the centre of which is O ; E is a fixed point without the circle; an ellipse is described with centre O and area constant so as always to touch EP at P . Find the locus of the extremities of the diameter conjugate to OP .

36. The normal at any point P of an ellipse cuts the axes in G, g ; prove that if any circle be described passing through G, g , the tangent to it from P is equal to CD .

37. Given a focus, a tangent, and the eccentricity* of a conic section; prove that the locus of the centre is a circle.

38. A straight line is drawn through a given point C within a circle to cut it in P, P' . If p is taken in it such that $Cp^2 = CP \cdot CP'$, find the locus of p .

39. In the ellipse $PY \cdot PY' : PN^2 :: CS^2 : BC^2$ and $SY \cdot CD = SP \cdot BC$.

40. Show that if the distance between the foci of the ellipse be greater than the length of its axis minor, there will be four positions of the tangent, for which the area of the triangle, included between it and the straight lines drawn from the centre of the curve to the feet of the perpendiculars from the foci on the tangent, will be the greatest possible.

* The ratio of $SA : AX$, or of $CS : CA$, is called the eccentricity.

41. Two conjugate diameters of an ellipse are cut by the tangent at any point P in M, N ; prove that the area of the triangle CPM varies inversely as that of the triangle CPN .

42. Circles are described on $SY, S'Y'$ as diameters, cutting $SP, S'P$ respectively in Q, Q' . Prove that $SQ \cdot S'P = SP \cdot S'Q' = BC^2$.

43. PSP', pSp' are any two focal chords of a conic section, P and p being on the same side of the axis; prove that $Pp, P'p'$ meet on the directrix.

44. Prove that an ellipse can be inscribed in any parallelogram so as to touch the middle points of the four sides; and show that this ellipse is the greatest of all inscribed ellipses.

45. If from any point on the exterior of two concentric, similar, and similarly placed ellipses, two tangents be drawn to the interior ellipse which also intersect the exterior; show that the distance between the points of intersection will be double of that between the points of contact.

46. The tangent at any point P in an ellipse, of which S and H are the foci, meets the axis major in T , and TQR bisects HP in Q , and meets SP in R ; prove that PR is one-fourth of the chord of curvature at P through S .

47. Prove that the distance between the two points on the circumference of an ellipse at which a given chord, not passing through the centre, subtends the greatest and least angles, is equal to the diameter which bisects that chord.

48. From any point on the auxiliary circle chords are drawn through the foci of an ellipse, and straight lines join the extremities of the chords with the extremity of the diameter passing through the point; prove that these lines will touch the ellipse.

49. A quadrilateral circumscribes an ellipse. Prove that either pair of opposite sides subtends supplementary angles at either focus.

50. Two tangents to an ellipse intersect at right angles; show that the straight line joining their point of intersection with the point of intersection of the normals at the points of contact passes through the centre.

51. P, Q are points in two confocal ellipses, at which the line joining the common foci subtends equal angles; prove that the tangents at P, Q are inclined at an angle which is equal to the angle subtended by PQ at either focus.

52. Tangents to an ellipse are drawn from any point in a circle through the foci; prove that the lines bisecting the angle between the tangents, all pass through a fixed point.

53. If the ordinate at P meet the auxiliary circle in Q , and CQ meet the ellipse in R , then CR is equal to the perpendicular on the tangent at P from C .

54. If P be a point such that $SP, S'P$ are perpendicular; prove that $CD^2 = 2 \cdot BC^2$.

55. If circles be escribed to the triangle SPS' opposite to the angles S and S' ; prove that the rectangle contained by their radii is equal to BC^2 .

56. The circle of curvature at any point P of an ellipse meets the focal distances in R, R' ; SU is a tangent to the circle.

Prove that $SU^2 : SP^2 :: 2 \cdot SP - 3 \cdot AC : AC$,

and if RR' passes through the centre of the circle of curvature, $CP = CS$. Determine the limits of possibility in both cases.

57. A straight line is drawn from the centre of an ellipse meeting the ellipse in P , the circle on the major axis in Q , and the tangent at the vertex in T . Prove that as CT approaches and ultimately coincides with the semi-major axis, PT and QT are ultimately in the duplicate ratio of the axes.

58. A straight line is drawn through the focus S of an ellipse meeting the two tangents at right angles to it in Y and Z , the diameter parallel to these tangents in L , and the directrix in M ; prove that

$$SL : SY :: SZ : SM.$$

59. If any equilateral triangle PQR be described in the auxiliary circle of an ellipse, and the ordinates to P, Q, R meet the ellipse in P', Q', R' ; the circles of curvature at P', Q', R' , meet in one point lying on the ellipse.

60. From a point T two tangents TP, TQ are drawn to an ellipse. Show that a circle with T as centre can be described so as to touch $SP, S'P, SQ, S'Q$.

61. If the normal at P meet the axis minor in g , and if the tangent at P meet the tangent at the vertex A in V ; show that

$$Sg : SC :: PV : VA.$$

62. If a circle passing through Y and Y' touch the major axis in Q , and that diameter of the circle which passes through Q meet the tangent in P ; show that $PQ = BC$. (See fig. Prop. XV.)

63. If PG the normal at P cut the major axis in G , and if DR, PN be the ordinates of D and P , prove that the triangles PGN, DRC are similar; and thence deduce that PG bears a constant ratio to CD .

64. The tangent at a point P of an ellipse meets the tangents at the vertices in V, V' . On VV' as diameter, a circle is described which intersects the ellipse in Q, Q' ; show that the ordinate of Q is to the ordinate of P as BC to $BC + CD$; where CD is conjugate to CP .

65. PCP' is any diameter of an ellipse; the tangents at any two points E and E' intersect in F ; $PE', P'E$ intersect in G . Show that FG is parallel to the diameter conjugate to PCP' .

66. If P be any point on an ellipse, and with P as centre and the semi-axis minor as radius a circle be described; prove that if PG be the normal, a circle described on CG as diameter will cut the first circle at right angles.

67. ABC is an isosceles triangle having $AB = AC$. BD, BE drawn on opposite sides of BC , and equally inclined to it, meet AC in D and E .

68. If an ellipse be described about BDC having its minor axis parallel to BC ; then AB will be a tangent to the ellipse.

69. If AQ be drawn from one of the vertices of an ellipse perpendicular to the tangent at any point P ; prove that the locus of the point of intersection of PS and QA produced will be a circle.

70. If Y, Y' be the feet of the perpendiculars from the foci of an ellipse on the tangent at P ; prove that the circle circumscribed about the triangle YNY' will pass through C .

71. Prove that the angle between the tangents to the auxiliary circle at Y, Y' , is the supplement of the angle SPS' .

72. P is any point on an ellipse; PM, PN perpendicular to the axes meet respectively, when produced, the circles described on the axes as diameter in the points Q, Q' . Show that QQ' passes through the centre.

73. Assuming that the greatest triangle which can be inscribed in a circle is equilateral, prove by the method of projection, that the greatest triangle which can be inscribed in an ellipse has one of its sides bisected by a diameter of the ellipse, and the others cut in points of bisection by the conjugate diameter.

74. PQ is a chord of an ellipse, normal at P , LCL' the diameter bisecting it. Show that PQ bisects the angle LPL' , and that $LP + L'P$ is constant.

75. A tangent to an ellipse at a point P intersects a fixed tangent in T ; if through the focus a line be drawn perpendicular to ST meeting the tangent to P in Q ; show that the locus of Q is a straight line touching the ellipse.

76. In an ellipse if a line be drawn through the focus making a constant angle with the tangent; prove that the locus of the point of intersection with the tangent is a circle.

77. Any chord PP' of an ellipse is produced to a point Q , such that $P'Q$ is equal to half the diameter parallel to PP' , and $QR'R$ is drawn through the centre to meet the ellipse in R, R' ; show that the area PCR is three times the area PCR' .

78. In an ellipse, L is the extremity of the latus rectum, and CD conjugate to CL . If a circle be described with centre C and passing through B , and a line be drawn through D parallel to the major axis, the portion of this line which lies within the circle will be equal to the latus rectum.

79. If P be any point in an ellipse, and K the point in which a normal at P intersects a line at right angles to it through S' , E the point of intersection of SP , and the diameter conjugate to CP , and if EK and CK be joined, each of the figures $SCKE$, $S'CEK$ will be a parallelogram.

80. If T be a point on the axis AA' produced, and PN the ordinate of the point where the tangent from T touches the ellipse; prove that

$$AN \cdot A'N : AT \cdot A'T :: CN : CT.$$

81. Given in an ellipse a focus and two tangents; prove that the locus of the other focus is a straight line.

82. A focus, a tangent, and the axis major being given, prove that the locus of the other focus is a circle.

83. A focus, a tangent, and the axis minor being given, prove that the locus of the other focus is a straight line.

84. An ellipse touches a fixed ellipse and has a common focus with it; if the major axis be fixed, the locus of the other focus is a circle; if the minor axis is fixed, the locus is an ellipse.

85. An ellipse and a parabola have a common focus. Prove that the ellipse either intersects the parabola in two points, and has two common tangents with it, or else does not cut it.

86. If in the ellipse a focus, a point, and the axis minor be given, the locus of the other focus is a parabola.

87. If at the extremities P, Q of any two diameters CP, CQ of an ellipse, two tangents Pp, Qq be drawn cutting each other in T , and the diameter produced in p and q , then the areas of the triangles TQp, TPq are equal.

88. If a straight line CN be drawn from the centre to bisect that chord of the circle of curvature at any point P of an ellipse, which is common to the ellipse and circle, and if it be produced to cut the ellipse in Q , and the tangent in T ; prove that $CP = CQ$, and that each is a mean proportional between CN and CT .

89. A parabola of given latus rectum is described touching symmetrically two conjugate diameters of an ellipse; find the locus of the focus.

90. An ellipse is described so as to touch the three sides of a triangle; prove that if one of its foci move along the circumference of a circle passing through two of the angular points of the triangle, the other will move along the circumference of another circle, passing through the same two angular points. Prove also that if one of these circles pass through the centre of the circle inscribed in the triangle, the two circles will coincide.

91. A triangle is described about an ellipse, so that the extremities of one of its sides lie in an ellipse, confocal with the given one; prove that the line bisecting the opposite angle passes through the pole of that side with respect to the outer ellipse.

92. Prove the following construction for a pair of tangents from any external point T to an ellipse of which the centre is C . Join CT ; let $TPCP'T$, a similar and similarly situated ellipse, be drawn, of which CT is a diameter, and P, P' its points of intersection with the given ellipse; TP, TP' will be tangents to the given ellipse.

93. The locus of the foci of all ellipses inscribed in the same parallelogram is a rectangular hyperbola. Prove this, and give a geometrical construction for finding the asymptotes.

94. AC is a fixed diameter of a circle, $ABCD$ a quadrilateral figure inscribed in the circle; prove that if the angles BAC, DAC be complementary, the locus of the intersection of BA, CD will be an hyperbola.

95. Prove that a circle can be described so as to touch the four straight lines drawn from the foci of an hyperbola to any two points on the same branch of the curve.

96. Any three diameters of an ellipse LL', MM', NN' , being taken, a circumscribing parallelogram $RTUV$ touches the ellipse at L, L', M, M' . Show that a conic section can be described through the points R, T, U, V, N, N' , which will be an hyperbola whose asymptotes are the lines forming in the ellipse the diameters conjugate to NN' and to the other common chord of the ellipse and hyperbola.

97. On opposite angles of any chord of a rectangular hyperbola are described equal segments of circles; show that the four points in which the circles to which these segments belong again meet the hyperbola, are the angular points of a parallelogram.

98. A triangle is inscribed in a rectangular hyperbola; prove that the circle described through the middle points of the sides of the triangle passes through the centre of the hyperbola.

99. ACB is an isosceles triangle; AB the base, and D any point in CB or CB produced: if BZ be drawn parallel to AD , meeting CA or CA produced in Z , prove that the middle point of DZ will be in an hyperbola whose asymptotes are CA , CB .

100. An ellipse and hyperbola are described so that the foci of each are at the extremities of the transverse axis of the other; prove that the tangents at their points of intersection meet the conjugate axis in points equidistant from the centre.

101. In a rectangular hyperbola, PK , PL are drawn at right angles to $A'P$, AP respectively, to meet the transverse axis in K and L ; prove that PK is equal to AP and KL to AA' , and the normal at P bisects KL .

102. In a rectangular hyperbola PO is a fixed diameter, Q any point on the curve; show that the angles QPO , QOP differ by a constant angle.

103. If the tangent at any point P of an hyperbola cut an asymptote in T , and if SP cut the same asymptote in Q , then $SQ = QT$.

104. If a given point be the focus of any hyperbola, passing through a given point and touching a given straight line, prove that the locus of the other focus is an arc of a fixed hyperbola.

105. If a series of conics be described having equal latera recta, the focus of a given parabola as their common focus, and the tangents to the parabola as their directrices; the common tangents to any two of the series will intersect in the directrix of the parabola.

106. At any P of an hyperbola a tangent is drawn, and PQ is taken on it in a constant ratio to CD ; prove that the locus of Q is an hyperbola.

107. If an hyperbola be described touching the four sides of a quadrilateral inscribed in a circle, and one focus lie on a circle, the other focus will also lie on this circle.

108. In an hyperbola, supposing the two asymptotes and one point of the curve to be given in position, show how to construct the curve; and find the position of the foci.

109. If A, D be two fixed points, and the angle PAD always exceed PDA by a given angle; find the locus of P , and the position of the transverse axis and asymptote.

110. From the middle point D of the base AB of the triangle ABC a straight line EDE' is drawn, making a given angle with AB , and the points E, E' are taken so that $ED = E'D = \frac{1}{2} AB$. If CA, CB take all possible positions consistent with the condition that the difference of the angles CAB, CBA is equal to EDA ; prove that the point C will trace out a rectangular hyperbola of which AB, EE' are conjugate diameters.

111. In the rectangular hyperbola, prove that the triangle, formed by the tangent at any point and its intercepts on the axes, is similar to the triangle formed by the straight line joining that point with the centre, and the abscissa and semi-ordinate of the point.

112. Tangents are drawn to an hyperbola, and the portion of each tangent intercepted by the asymptotes is divided in a constant ratio; prove that the locus of the point of section is an hyperbola.

113. Show that the point of trisection of a series of conterminous circular arcs lie on branches of two hyperbolas and determine the distance between their centres.

114. From a point R on one asymptote RE is drawn touching the hyperbola in E , and ET , EV are drawn through E , parallel to the asymptotes, cutting a diameter in T and V ; RV is joined, cutting the hyperbola in P , p : show that TP , Tp touch the hyperbola.

115. Given in the ellipse a focus and two points, the locus of the other focus is an hyperbola.

116. If a rectangular hyperbola passes through three given points, the locus of its centre is a circle, which passes through the middle points of the lines joining the three given points.

117. If the tangent at P meet one asymptote in T , and a line TQ be drawn parallel to the other asymptote to meet the curve in Q ; prove that if PQ be joined and produced both ways to meet the asymptotes in R and R' , RR' will be trisected at the points P and Q .

118. If two concentric rectangular hyperbolas have a common tangent, the lines joining their points of intersection to their respective points of contact with the common tangent will subtend equal angles at their common centres.

119. If TP , TQ be two tangents drawn from any point T to touch a conic in P and Q , and if S and S' be the foci, then

$$ST^2 : S'T^2 :: SP \cdot SQ : S'P \cdot S'Q.$$

120. The circle of curvature at the vertex of a conic meets the axis again in D , and a tangent is drawn to the circle at D : if two tangents be drawn to the circle from any point in the conic they will intercept between them a constant length of the former tangent.

121. A family of hyperbolas is described about a given triangle; prove that if one of the asymptotes always passes through a fixed point the other will always touch a fixed conic section to which the three sides of the triangle are tangents.

122. If the lines which bisect the angles between pairs of tangents to an ellipse be parallel to a fixed straight line, prove that the locus of the points of intersection of the tangents will be a rectangular hyperbola.

123. An hyperbola, of given eccentricity, always passes through two given points; if one of its asymptotes always pass through a third given point in the same straight line with these, prove that the locus of the centre of the hyperbola will be a circle.

124. A, P and B, Q are points taken respectively in two parallel straight lines, A and B being fixed, and P, Q variable. Prove that if the rectangle $APBQ$ be constant, the line PQ will always touch a fixed ellipse or a fixed hyperbola, according as P and Q are on the same or opposite sides of AB .

125. If two plane sections of a right cone be taken, having the same directrix, the foci corresponding to that directrix lie on a straight line which passes through the vertex.

126. Give a geometrical construction by which a cone may be cut so that the section may be an ellipse of given eccentricity.

127. Given a right cone and a point within it, there are but two sections which have this point for focus; and the planes of these sections make equal angles with the straight line joining the given point and the vertex of the cone.

128. If the curve formed by the intersection of any plane with a cone be projected upon a plane perpendicular to the axis, prove that the curve of projection will be a conic section, having its focus at the point in which the axis meets the plane of projection.

129. If F be the point where the major axis of an elliptic

section meets the axis of the cone, and C be the centre of the section ; prove that

$$CF : CS :: AA' : AO + A'O,$$

O being the vertex of the cone.

130. Two intersecting right cones have the same axis, and the vertical angle of the interior is equal to that of an equilateral triangle ; show that the vertex of the interior cone is a focus of all sections of the exterior made by planes touching the interior.

THE END.

London and Cambridge.

MACMILLAN AND CO.'S
DESCRIPTIVE CATALOGUE OF
CAMBRIDGE CLASS BOOKS.

The Works in this Series of CAMBRIDGE CLASS-BOOKS FOR THE USE OF SCHOOLS AND COLLEGES, which have been issued at intervals during the last ten years, are intended to embrace all branches of Education, from the most Elementary to the most Advanced, and to keep pace with the latest discoveries in Science.

Of those hitherto published the large and ever increasing sale is a sufficient indication of the manner in which they have been appreciated by the public.

A SERIES of a more Elementary character is in preparation, a list of which will be found on page 2 of this Catalogue.

ELEMENTARY SCHOOL CLASS BOOKS.

The volumes of this Series of ELEMENTARY SCHOOL CLASS BOOKS are handsomely printed in a form that, it is hoped, will assist the young student as much as clearness of type and distinctness of arrangement can effect. They are published at a moderate price to ensure an extensive sale in the Schools of the United Kingdom and the Colonies.

1. *Euclid for Colleges and Schools.*

By I. TODD HUNTER, M.A., F.R.S., Fellow and Principal Mathematical Lecturer of St. John's College, Cambridge. 18mo. 3s. 6d.

2. *An Elementary Latin Grammar.*

By H. J. ROBY, M.A., Under Master of Dulwich College Upper School, late Fellow and Classical Lecturer of St. John's College, Cambridge. 18mo. 2s. 6d.

3. *An Elementary History of the Book of Common Prayer.*

By FRANCIS PROCTER, M.A., Vicar of Wotton, Norfolk, late Fellow of St. Catharine's College, Cambridge. 18mo. 2s. 6d.

4. *Mythology for Latin Versification.*

A Brief Sketch of the Fables of the Ancients, prepared to be rendered into Latin Verse, for Schools. By F. C. HODGSON, B.D., late Provost of Eton College. *New Edition.* Revised by F. C. HODGSON, M.A., Fellow of King's College, Cambridge. 18mo. 3s.

5. *Algebra for Beginners.*

By I. TODD HUNTER, M.A., F.R.S. [In the Press.

6. *The School Class Book of Arithmetic.*

By BARNARD SMITH, M.A., late Fellow of St. Peter's College, Cambridge. [In the Press.

7. *The Bible Word-Book.*

A Glossary of old English Bible Words with Illustrations. By J. EASTWOOD, M.A., St. John's College, Cambridge, and Incumbent of Hope in Hanley, Stafford, and W. ALDIS WRIGHT, M.A., Trinity College, Cambridge. [Preparing.

* * * Other Volumes will be announced in due course.

CAMBRIDGE CLASS BOOKS

FOR

SCHOOLS AND COLLEGES.

WORKS by the Rev. BARNARD SMITH, M.A.

Fellow of St. Peter's College, Cambridge. •

1.

Arithmetic & Algebra

In their Principles and Applications.

With numerous Examples, systematically arranged. •

Eighth Edit. 696 pp. (1861). Cr. 8vo. strongly bound in cloth. 10s. 6d.

The first edition of this work was published in 1854. It was primarily intended for the use of students at the Universities, and for Schools which prepare for the Universities. It has however been found to meet the requirements of a much larger class, and is now extensively used in *Schools and Colleges both at home and in the Colonies*. It has also been found of great service for students preparing for the MIDDLE-CLASS and CIVIL AND MILITARY SERVICE EXAMINATIONS, from the care that has been taken to elucidate the *principles* of all the Rules. Testimony of its excellence has been borne by some of the highest practical and theoretical authorities, of which the following from the late DEAN PEACOCK may be taken as a specimen:

"Mr. Smith's Work is a most useful publication. The Rules are stated with great clearness. The Examples are well selected and worked out with just sufficient detail without being encumbered by too minute explanations; and there prevails throughout it that just proportion of theory and practice, which is the crowning excellence of an elementary work."

2. **Arithmetic**

For the Use of Schools.

New Edition (1862) 348 pp.

Crown 8vo strongly bound in cloth, 4s. 6d. Answers to all the Questions.

3. **Key** to the above, containing Solutions to all the Questions in the latest Edition. Crown 8vo. cloth. 392 pp. Second Edit. 8s. 6d.

To meet a widely expressed wish, the ARITHMETIC was published separately from the larger work in 1854, with so much alteration as was necessary to make it quite independent of the ALGEBRA. It has now a very large sale in all classes of schools at home and in the Colonies. A copious collection of Examples, under each rule, has been embodied in the work in a systematic order, and a Collection of Miscellaneous Papers in all branches of Arithmetic is appended to the book.

4. **Exercises in Arithmetic.** 104 pp. Cr. 8vo. (1860) 2s. Or with ANSWERS, 2s. 6d. Also sold separately in 2 Parts 1s. each. **Answers, 6d.**

These EXERCISES have been published in order to give the pupil examples in every rule of Arithmetic. The greater number have been carefully compiled from the latest University and School Examination Papers.

CAMBRIDGE CLASS BOOKS

WORKS by ISAAC TODHUNTER, M.A. F.R.S.

Fellow and Principal Mathematical Lecturer of St. John's College, Cambridge.

1. Algebra.

For the Use of Colleges and Schools.

Third Edition. 542 pp. (1862).
Strongly bound in cloth. 7s. 6d.

This work contains all the propositions which are usually included in elementary treatises on Algebra, and a large number of *Examples for Exercise*. The author has sought to render the work easily intelligible to students without impairing the accuracy of the demonstrations, or contracting the limits of the subject. The Examples have been selected with a view to illustrate every part of the subject, and as the number of them is about *Sixteen hundred and fifty*, it is hoped they will supply ample exercise for the student. Each set of Examples has been carefully arranged, commencing with very simple exercises, and proceeding gradually to those which are less obvious.

2. Plane Trigonometry

For Schools and Colleges.

2nd Edit. 279 pp. (1860). Crn. 8vo.
Strongly bound in cloth. 5s.

The design of this work has been to render the subject intelligible to beginners, and at the same time to afford the student the opportunity of obtaining all the information which he will require on this branch of Mathematics. Each chapter is followed by a set of Examples; those which are entitled *Miscellaneous Examples*, together with a few in some of the other sets, may be advantageously reserved by the student for exercise after he has made some progress in the subject. As the Text and Examples have been tested by considerable experience in teaching, the hope is entertained that they will be suitable for imparting a sound and comprehensive knowledge of Plane Trigonometry, together with readiness in the application of this knowledge to the solution of problems. In the Second Edition the hints for the solution of the Examples have been considerably increased.

3. Spherical Trigonometry.

For the Use of Colleges and Schools.

112 pp. Crown 8vo. (1859).
Strongly bound in cloth. 4s. 6d.

This work is constructed on the same plan as the *Treatise on Plane Trigonometry*, to which it is intended as a sequel. Considerable labour has been expended on the text in order to render it comprehensive and accurate, and the Examples, which have been chiefly selected from University and College Papers, have all been carefully verified.

The Elements of Euclid

For the Use of Schools and Colleges

COMPRISING THE FIRST SIX BOOKS AND PORTIONS OF THE ELEVENTH AND TWELFTH BOOKS, WITH NOTES, APPENDIX, AND EXERCISES.

384 pp. 18mo. bound. (1862). 3s. 6d

As the Elements of Euclid are usually placed in the hands of young students, it is important to exhibit the work in such a form as will assist them in overcoming the difficulties which they experience on their first introduction to processes of continuous argument. No method appears to be so useful as that of breaking up the demonstrations into their constituent parts, and this plan has been adopted in the present edition. Each distinct assertion in the argument begins a new line, and at the end of the lines are placed the necessary references to the preceding principles on which the assertions depend. The longer propositions are distributed into subordinate parts, which are distinguished by breaks at the beginning of the lines. The Notes are intended to indicate and explain the principal difficulties, and to supply the most important inferences which can be drawn from the propositions. The work finishes with a collection of *Six hundred and twenty-five Exercises*, which have been selected principally from Cambridge Examination papers and have been tested by long experience. As far as possible they are arranged in order of difficulty. The Figures will be found to be large and distinct, and have been repeated when necessary, so that they always occur in immediate connexion with the corresponding text.

WORKS by ISAAC TODHUNTER, M.A., F.R.S.—*continued.*

. . 5.

The Integral Calculus*And its Applications.*

With numerous Examples.

Second Edition. 342 pp. (1862).

Crown 8vo. cloth. 10s. 6d.

In writing the present *Treatise on the Integral Calculus*, the object has been to produce a work at once elementary and complete—adapted for the use of beginners, and sufficient for the wants of advanced students. In the selection of the propositions, and in the mode of establishing them, the author has endeavoured to exhibit fully and clearly the principles of the subject, and to illustrate all their most important results. In order that the student may find in the volume all that he requires, a large collection of Examples for exercise has been appended to the different chapters.

6. Analytical Statics.*With numerous Examples.*

Second Edition. 230 pp. (1858).

Crown 8vo. cloth. 10s. 6d.

In this work will be found all the propositions which usually appear in treatises on Theoretical Statics. To the different chapters Examples are appended, which have been selected principally from the University and College Examination Papers; these will furnish ample exercise in the application of the principles of the subject.

7. EXAMPLES OF**Analytical Geometry of Three Dimensions.**

76 pp. (1858). Crn. 8vo. cloth. 4s.

A collection of examples in illustration of Analytical Geometry of Three Dimensions has long been required both by students and teachers, and the present work is published with the view of supplying the want.

8. The**Differential Calculus.***With numerous Examples.*

Third Edition, 398 pp. (1860).

Crown 8vo. cloth, 10s. 6d.

This work is intended to exhibit a comprehensive view of the Differential Calculus on the method of Limits. In the more elementary portions, explanations have been given in considerable detail, with the hope that a reader who is without the assistance of a tutor may be enabled to acquire a competent acquaintance with the subject. More than one investigation of a theorem has been frequently given, because it is believed that the student derives advantage from viewing the same proposition under different aspects, and that in order to succeed in the examinations which he may have to undergo, he should be prepared for a considerable variety in the order of arranging the several branches of the subject, and for a corresponding variety in the mode of demonstration.

9. Plane Co-Ordinate Geometry

AS APPLIED TO THE STRAIGHT LINE AND THE CONIC SECTIONS.

With numerous Examples.

Third and Cheaper Edition.

Crn. 8vo. cl. 326 pp. (1862). 7s. 6d.

This *Treatise* exhibits the subject in a simple manner for the benefit of beginners, and at the same time includes in one volume all that students usually require. The Examples at the end of each chapter will, it is hoped, furnish sufficient exercise, as they have been carefully selected with the view of illustrating the most important points, and have been tested by repeated experience with pupils. In consequence of the demand for the work proving much greater than had been originally anticipated, a large number of copies of the *Third Edition* has been printed, and a considerable reduction effected in the price.

By ISAAC TODHUNTER, M.A.

10. **ELEMENTARY TREATISE ON THE
Theory of Equations.**

With a Collection of Examples.

Crown 8vo. cloth. 279 pp. (1861).
7s. 6d.

This treatise contains all the propositions which are usually included in elementary treatises on the Theory of Equations, together with a collection of Examples for exercise. This work may in fact be regarded as a sequel to that on Algebra by the same writer, and accordingly the student has occasionally been referred to the treatise on Algebra for preliminary information on some topics here discussed. The work includes three chapters on Determinants.

11. **History of the Progress
of the
Calculus of Variations**

During the Nineteenth Century.

8vo. cloth. 532 pp. (1861). 12s.

It is of importance that those who wish to cultivate any subject may be able to ascertain what results have already been obtained, and thus reserve their strength for difficulties which have not yet been conquered. The Author has endeavoured in this work to ascertain distinctly what has been effected in the Progress of the Calculus, and to form some estimate of the manner in which it has been effected.



A TREATISE ON
**Mechanics and Hydro-
statics.**

With Solutions of Questions

PROPOSED IN THE CAMBRIDGE SENATE HOUSE

By W. H. GIRDLESTONE, M.A.

Christ's College.

8vo. cloth. 100 pp. 1862.

By J. H. PRATT, M.A.

Archdeacon of Calcutta, late Fellow of
Gonville and Caius College, Cambridge.

**A Treatise on
Attractions,**

*La Place's Functions, and the Figure
of the Earth.*

Second Edition. Crown 8vo. 126 pp.
(1861). cloth. 6s. 6d.

In the present Treatise the author has endeavoured to supply the want of a work on a subject of great importance and high interest—La Place's Coefficients and Functions and the calculation of the Figure of the Earth by means of his remarkable analysis. No student of the higher branches of Physical Astronomy should be ignorant of La Place's analysis and its result—"a calculus," says Argy, "the most singular in its nature and the most powerful in its application that has ever appeared."



By G. B. AIRY, M.A.

Astronomer Royal.

1. **Mathematical Tracts**

*On the Lunar and Planetary Theories,
Figure of the Earth, the Undulatory
Theory of Optics, &c.*

Fourth Edition. 400 pp. (1858).
8vo. 15s.

2. **Theory of Errors of
Observations**

And the Combination of Observations.
103 pp. (1861). Crown 8vo. 6s. 6d.

In order to spare astronomers and observers in natural philosophy the confusion and loss of time which are produced by referring to the ordinary treatises embracing both branches of Probabilities, the author has thought it desirable to draw up this work, relating only to Errors of Observation, and to the rules derivable from the consideration of these Errors, for the Combination of the Results of Observations. The Author has thus also the advantage of entering somewhat more fully into several points of interest to the observer, than can possibly be done in a General Theory of Probabilities.

By **GEORGE BOOLE, D.C.L., F.R.S.**
Professor of Mathematics in the Queen's
University, Ireland.

Differential Equations

468 pp. (1859). Crn. 8vo. cloth. 14s.

The Author has endeavoured in this treatise to convey as complete an account of the present state of knowledge on the subject of Differential Equations as was consistent with the idea of a work intended, primarily, for elementary instruction. The object has been first of all to meet the wants of those who had no previous acquaintance with the subject, and also not quite to disappoint others who might seek for more advanced information. The earlier sections of each chapter contain that kind of matter which has usually been thought suitable for the beginner, while the latter ones are devoted either to an account of recent discovery, or to the discussion of such deeper questions of principle as are likely to present themselves to the reflective student in connection with the methods and processes of his previous course.

2. The Calculus of Finite Differences.

248 pp. (1860). Crown 8vo. cloth.
10s. 6d.

In this work particular attention has been paid to the connexion of the methods with those of the Differential Calculus—a connexion which in some instances involves far more than a merely formal analogy. The work is in some measure designed as a sequel to the Author's *Treatise on Differential Equations*, and it has been composed on the same plan.

Elementary Statics.

By the *Rev. GEORGE RAWLINSON*
Professor of Applied Sciences, Elphinstone Coll., Bombay.

Edited by the Rev. E. STURGES, M.A.
Rector of Kencott, Oxfordshire.

(150 pp.) 1860. Crn. 8vo. cl. 4s. 6d.

This work is published under the authority of H. M. Secretary of State for India for use in the Government Schools and Colleges in India.

By **P. G. TAIT, M.A.**, and
W. J. STEELE, B.A.

Late Fellows of St. Peter's Coll. Camb.

Dynamics of a Particle.

With numerous Examples.

304 pp. (1856). Cr. 8vo. cl. 10s. 6d.

In this Treatise will be found all the ordinary propositions connected with the Dynamics of Particles which can be conveniently deduced without the use of D'Alembert's Principles. Throughout the book will be found a number of illustrative Examples introduced in the text, and for the most part completely worked out; others, with occasional solutions or hints to assist the student are appended to each Chapter.

By the *Rev. G. F. CHILDE, M.A.*
Mathematical Professor in the South
African College.

Singular Properties of the Ellipsoid

And Associated Surfaces of the n^{th} Degree.

152 pp. (1861). 8vo. boards. 10s. 6d.

As the title of this volume indicates, its object is to develop peculiarities in the Ellipsoid; and further, to establish analogous properties in unlimited congeneric series of which this remarkable surface is a constituent.

By **J. B. PHEAR, M.A.**
Fellow and late Mathematical Lecturer of
Clare College.

Elementary Hydrostatics

With numerous Examples and Solutions.

Second Edition. 156 pp. (1857)
Crown 8vo. cloth. 5s. 6d.

"An excellent Introductory Book. The definitions are very clear; the descriptions and explanations are sufficiently full and intelligible; the investigations are simple and scientific. The examples greatly enhance its value."—*ENGLISH JOURNAL OF EDUCATION*.

This Edition contains 117 Examples, and solutions to all these examples are given at the end of the book.

By *Rev. S. PARKINSON, B.D.*

Fellow and Praelector of St. John's Coll.
Cambridge.

1. **Elementary Treatise on Mechanics.**

With a Collection of Examples.

Second Edition. 345 pp. (1861).
Crown 8vo. cloth. 9s. 6d.

The Author has endeavoured to render the present volume suitable as a Manual for the junior classes in Universities and the higher classes in Schools. With this object there have been included in it those portions of theoretical Mechanics which can be conveniently investigated without the Differential Calculus, and with one or two short exceptions the student is not presumed to require a knowledge of any branches of Mathematics beyond the elements of Algebra, Geometry, and Trigonometry. A collection of Problems and Examples has been added, chiefly taken from the Senate-House and College Examination Papers—which will be found useful as an exercise for the student. In the Second Edition several additional propositions have been incorporated in the work for the purpose of rendering it more complete, and the Collection of Examples and Problems has been largely increased.

2. **A Treatise on Optics**

304 pp. (1859). Crown 8vo. 10s. 6d.

A collection of Examples and Problems has been appended to this work which are sufficiently numerous and varied in character to afford useful exercise for the student: for the greater part of them recourse has been had to the Examination Papers set in the University and the several Colleges during the last twenty years.

Subjoined to the copious Table of Contents the author has ventured to indicate an elementary course of reading not unsuitable for the requirements of the First Three Days in the Cambridge Senate House Examinations.

By *R. D. BEASLEY, M.A.*

Head Master of Grantham School.

AN ELEMENTARY TREATISE ON **Plane Trigonometry.**

*With a numerous Collection of
Examples.*

106 pp. (1858), strongly bound in
cloth. 3s. 6d.

This Treatise is specially intended for use in Schools. The choice of matter has been chiefly guided by the requirements of the three days' Examination at Cambridge, with the exception of proportional parts in logarithms, which have been omitted. About *Four hundred* Examples have been added, mainly collected from the Examination Papers of the last ten years, and great pains have been taken to exclude from the body of the work any which might dishearten a beginner by their difficulty.

By *J. BROOK SMITH, M.A.*

St. John's College, Cambridge.

Arithmetic in Theory and Practice.

For Advanced Pupils.

PART I. Crown 8vo. cloth. 3s. 6d.

This work forms the first part of a Treatise on Arithmetic, in which the Author has endeavoured, from very simple principles, to explain in a full and satisfactory manner all the important processes in that subject.

The proofs have in all cases been given in a form entirely arithmetical: for the author does not think that recourse ought to be had to Algebra until the arithmetical proof has become hopelessly long and perplexing.

At the end of every chapter several examples have been worked out at length, in which the best practical methods of operation have been carefully pointed out.

By G. H. PUCKLE, M.A.

Principal of Windermere College.

Conic Sections and Algebraic Geometry.

With numerous Easy Examples Progressively arranged.

Second Edition. 264 pp. (1856).
Crown 8vo. 7s. 6d.

This book has been written with special reference to those difficulties and misapprehensions which commonly beset the student when he commences. With this object in view, the earlier part of the subject has been dwelt on at length, and geometrical and numerical illustrations of the analysis have been introduced. The Examples appended to each section are mostly of an elementary description. The work will, it is hoped, be found to contain all that is required by the upper classes of schools and by the generality of students at the Universities.

By EDWARD JOHN ROUTH, M.A.

Fellow and Assistant Tutor of St. Peter's College, Cambridge.

Dynamics of a System of Rigid Bodies.

With numerous Examples.

336 pp. (1860). Crown 8vo. cloth.
10s. 6d.

CONTENTS: Chap. I. Of Moments of Inertia.—II. D'Alembert's Principle.—III. Motion about a Fixed Axis.—IV. Motion in Two Dimensions.—V. Motion of a Rigid Body in Three Dimensions.—VI. Motion of a Flexible String.—VII. Motion of a System of Rigid Bodies.—VIII. Of Impulsive Forces.—IX. Miscellaneous Examples.

The numerous Examples which will be found at the end of each chapter have been chiefly selected from the Examination Papers set in the University and Colleges of Cambridge during the last few years.

The Cambridge Year Book AND UNIVERSITY ALMANACK

For 1863.

Crown 8vo. 228 pp. price 2s. 6d.

The specific features of this annual publication will be obvious at a glance, and its value to teachers engaged in preparing students for, and to parents who are sending their sons to, the University, and to the public generally, will be clear.

1. The whole mode of proceeding in entering a student at the University and at any particular College is stated.

2. The course of the studies as regulated by the University examinations, the manner of these examinations, and the specific subjects and times for the year 1863, are given.

3. A complete account of all Scholarships and Exhibitions at the several Colleges, their value, and the means by which they are gained.

4. A brief summary of all Graces of the Senate, Degrees conferred during the year 1861, and University news generally are given.

5. The Regulations for the Local Examination of those who are not members of the University, to be held this year, with the names of the books on which the Examination will be based, and the date on which the Examination will be held.

By N. M. FERRERS, M.A.

Fellow and Mathematical Lecturer of Gonville and Caius College, Cambridge.

AN ELEMENTARY TREATISE ON Trilinear Co-Ordinates

*The Method of Reciprocal Polars,
and the Theory of Projections.*

154 pp. (1861). Cr. 8vo. cl. 6s. 6d.

The object of the Author in writing on this subject has mainly been to place it on a basis altogether independent of the ordinary Cartesian System, instead of regarding it as only a special form of abridged Notation. A short chapter on Determinants has been introduced.

By J. C. SNOWBALL, M.A.

Late Fellow of St. John's Coll. Cambridge.

Plane and Spherical Trigonometry.

With the Construction and Use of Tables of Logarithms.

Ninth Edition. 240 pp. (1857).
Crown 8vo. 7s. 6d.

In preparing a new edition, the proofs of some of the more important propositions have been rendered more strict and general, and a considerable addition of more than *Two hundred Examples*, taken principally from the questions in the Examinations of Colleges and the University, has been made to the collection of Examples and Problems for practice.

By W. H. DREW, M.A.

Second Master of Blackheath School.

Geometrical Treatise on Conic Sections.

With a copious Collection of Examples.

Second Edition. Crown 8vo. cloth.
4s. 6d.

In this work the subject of Conic Sections has been placed before the student in such a form that, it is hoped, after mastering the elements of Euclid, he may find it an easy and interesting continuation of his geometrical studies. With a view also of rendering the work a complete Manual of what is required at the Universities, there have been either embodied into the text, or inserted among the examples, every book work question, problem, and rider, which has been proposed in the Cambridge examinations up to the present time.

Solutions to the Problems in Drew's Conic Sections.

Crown 8vo. cloth. 4s. 6d.

Senate-House Mathematical Problems.

With Solutions.

1848-51. By FERRERS and JACKSON. 8vo.
15s. 6d.

1848-51. (RIDERS) By JAMISON. 8vo.
7s. 6d.

1854. By WALTON and MACKENZIE.
10s. 6d.

1857. By CAMPION and WALTON. 8vo.
8s. 6d.

1860. By ROUTH and WATSON. Crown
8vo. 7s. 6d.

The above books contain Problems and Examples which have been set in the Cambridge Senate-house Examinations at various periods during the last twelve years, together with Solutions of the same. The Solutions are in all cases given by the Examiners themselves or under their sanction.

— ♦ —

By H. A. MORGAN, M.A.

Fellow of Jesus College, Cambridge.

A Collection of Mathematical Problems and Examples.

With Answers.

190 pp. (1858). Crown 8vo. 6s. 6d.

This book contains a number of problems, chiefly elementary, in the Mathematical subjects usually read at Cambridge. They have been selected from the papers set during late years at Jesus College. Very few of them are to be met with in other collections, and by far the larger number are due to some of the most distinguished Mathematicians in the University.

— ♦ —

Cambridge University Examination Papers.

Crown 8vo. 184 pp. 2s. 6d.

A Collection of all the Papers set at the Examinations for the Degrees, the various Triposes, and the Theological Certificates in the University, with List of Candidates Examined and of those Approved, and an Index to the Subjects. 1860-61.

A Treatise on Solid Geometry.

By *PERCIVAL FROST, M.A.*,
St. John's College, and
JOSEPH WOLSTENHOLME, M.A.,
Christ's Coll. Cambridge.

472 pp. 8vo. cloth. 18s. 1863.

The authors have endeavoured to present before students as comprehensive a view of the subject as possible. Intending as they have done to make the subject accessible, at least in the earlier portion, to all classes of students, they have endeavoured to explain fully all the processes which are most useful in dealing with ordinary theorems and problems, thus directing the student to the selection of methods which are best adapted to the exigencies of each problem. In the more difficult portions of the subject, they have considered themselves to be addressing a higher class of students, there they have tried to lay a good foundation on which to build, if any reader should wish to pursue the science beyond the limits to which the work extends.



AN ELEMENTARY TREATISE ON The Planetary Theory.

WITH A COLLECTION OF PROBLEMS.

By *C. H. H. CHEYNE, B.A.*

Scholar of St. John's College, Cambridge.

148 pp. 1862. Crn. 8vo. cloth. 6s. 6d.

In this volume, an attempt has been made to produce a Treatise on the Planetary Theory, which being elementary in character, should be so far complete, as to contain all that is usually required by students in the University. A collection of Problems has been added, taken chiefly from Cambridge Examination papers of the last twenty years.

By *JOHN E. B. MAYOR, M.A.*

Fellow and Classical Lecturer of St. John's College, Cambridge.

1. Juvenal.

With English Notes.

464 pp. (1854). Crown 8vo. cloth.
10s. 6d.

"A School edition of Juvenal, which, for really ripe scholarship, extensive acquaintance with Latin literature, and familiar knowledge of Continental criticism, ancient and modern, is unsurpassed, we do not say among English School-books, but among English editions generally"—
EDINBURGH REVIEW.



2. Cicero's Second Philippic.

With English Notes.

168 pp. (1861). Fcp. 8vo. cloth. 5s.

The Text is that of Halm's 2nd edition, (Leipzig, Weidmann, 1858), with some corrections from Madvig's 4th Edition (Copenhagen, 1858). Halm's Introduction has been closely translated, with some additions. His notes have been curtailed, omitted, or enlarged, at discretion; passages to which he gives a bare reference, are for the most part printed at length; for the Greek extracts an English version has been substituted. A large body of notes, chiefly grammatical and historical, has been added from various sources. A list of books useful to the student of Cicero, a copious Argument, and an Index to the introduction and notes, complete the book.



By *P. FROST, Jun., M.A.*

Late Fellow of St. John's Coll. Cambridge.

Thucydides. Book VI.

With English Notes, Map and Index.

8vo. cloth. 7s. 6d.

It has been attempted in this work to facilitate the attainment of accuracy in translation. With this end in view the Text has been treated grammatically.

By B. DRAKE, M.A.

Late Fellow of King's Coll. Cambridge.

1. Demosthenes on the Crown.

With English Notes.

Second Edition. To which is prefixed ÆSCHINES AGAINST CTESIPHON. With English Notes.

287 pp. (1860). Fcap. 8vo. cl. 5s.

The first edition of the late Mr. Drake's edition of Demosthenes de Corona having met with considerable acceptance in various Schools, and a new edition being called for, the Oration of Æschines against Ctesiphon, in accordance with the wishes of many teachers, has been appended with useful notes by a competent scholar.

2. Æschyli Eumenides

With English Verse Translation, Copious Introduction, and Notes.

8vo. 144. pp. (1853). 7s. 6d.

"Mr. Drake's ability as a critical Scholar is known and admitted. In the edition of the Eumenides before us we meet with him also in the capacity of a Poet and Historical Essayist. The translation is flowing and melodious, elegant and scholarlike. The Greek Text is well printed: the notes are clear and useful"—GLARDIAN.

By C. MERIVALE, B.D.

Author of "History of Rome," &c.

Sallust.

With English Notes.

Second Edition. 172 pp. (1858).

Fcap. 8vo. 4s. 6d.

"This School edition of Sallust is precisely what the School edition of a Latin author ought to be. No useless words are spent in it, and no words that could be of use are spared. The text has been carefully collated with the best editions. With the work is given a full current of extremely well-selected annotations."—THE EXAMINER.

The "CATILINA" and "JUGURTHA" may be had separately, price 2s. 6d. each, bound in cloth.

By J. WRIGHT, M.A.

Head Master of Sutton Coldfield School

1. Help to Latin Grammar.

With Easy Exercises, and Vocabulary

Crown 8vo. cloth. 4s. 6d.

Never was there a better aid offered alike to teacher and scholar in that arduous pass. The style is at once familiar and strikingly simple and lucid; and the explanations precisely hit the difficulties, and thoroughly explain them."—ENGLISH JOURNAL OF EDUCATION.

2. Hellenica.

A FIRST GREEK READING BOOK.

Second Edit. Fcap. 8vo. cl. 3s. 6d.

In the last twenty chapters of this volume, Thucydides sketches the rise and progress of the Athenian Empire in so clear a style and in such simple language, that the author doubts whether any easier or more instructive passages can be selected for the use of the pupil who is commencing Greek.

3. The Seven Kings of Rome.

A First Latin Reading Book.

Second Edit. Fcap. 8vo. cloth. 3s.

This work is intended to supply the pupil with an easy Construing-book, which may, at the same time, be made the vehicle for instructing him in the rules of grammar and principles of composition. Here Livy tells his own pleasant stories in his own pleasant words. Let Livy be the master to teach a boy Latin, not some English collector of sentences, and he will not be found a dull one.

4. Vocabulary and Exercises on "The Seven Kings of Rome."

Fcap. 8vo. cloth. 2s. 6d.

* * * The Vocabulary and Exercises may also be had bound up with "The Seven Kings of Rome." 5s. cloth.

By EDWARD THRING, M.A.

Head Master of Uppingham School.

Elements of Grammar Taught in English.

With Questions.

Third Edition. 136 pp. (1860).
Demy 18mo. 2s.

2. The Child's English Grammar.

New Edition. 86 pp. (1859). Demy
18mo. 1s.

The Author's effort in these two books has been to point out the broad, beaten, every-day path, carefully avoiding digressions into the byeways and eccentricities of language. This Work took its rise from questionings in National Schools, and the whole of the first part is merely the writing out in order the answers to questions which have been used already with success. The study of Grammar in English has been much neglected, nay by some put on one side as an impossibility. There was perhaps much ground for this opinion, in the medley of arbitrary rules thrown before the student, which applied indeed to a certain number of instances, but would not work at all in many others, as must always be the case when principles are not put forward in a language full of ambiguities. The present work does not, therefore, pretend to be a compendium of idioms, or a philological treatise, but a Grammar. Or in other words, its intention is to teach the learner how to speak and write correctly, and to understand and explain the speech and writings of others. Its success, not only in National Schools, from practical work in which it took its rise, but also in classical schools, is full of encouragement.

3. School Songs.

A COLLECTION OF SONGS FOR
SCHOOLS.

WITH THE MUSIC ARRANGED FOR
FOUR VOICES.

*Edited by Rev E. THRING and
H. RICCIUS.*

Music Size. 7s. 6d.

By EDWARD THRING, M.A.

4. A First Latin Con- struing Book.

104 pp. (1855). Fcap. 8vo. 2s. 6d.

This Construing Book is drawn up on the same sort of graduated scale as the Author's *English Grammar*. Passages out of the best Latin Poets are gradually built up into their perfect shape. The few words altered, or inserted as the passages go on, are printed in Italics. It is hoped by this plan that the learner, whilst acquiring the rudiments of language, may store his mind with good poetry and a good vocabulary.

By C. J. VAUGHAN, D.D.

Head Master of Harrow School.

St. Paul's Epistle to the Romans.

The Greek Text with English Notes.

Second Edition. Crown 8vo. cloth.
(1861). 5s.

By dedicating this work to his elder Pupils at Harrow, the Author hopes that he sufficiently indicates what is and what is not to be looked for in it. He desires to record his impression, derived from the experience of many years, that the Epistles of the New Testament, no less than the Gospels, are capable of furnishing useful and solid instruction to the highest classes of our Public Schools. If they are taught accurately, not controversially; positively, not negatively; authoritatively, yet not dogmatically; taught with close and constant reference to their literal meaning, to the connexion of their parts, to the sequence of their argument, as well as to their moral and spiritual instruction; they will interest, they will inform, they will elevate; they will inspire a reverence for Scripture never to be discarded, they will awaken a desire to drink more deeply of the Word of God, certain hereafter to be gratified and fulfilled.

1.

By C. J. VAUGHAN, D.D.

**Notes for
Lectures on Confirmation.**
With Suitable Prayers.
4th Edition. 70 pp. (1862). Fcp.
8vo. 1s. 6d.

This work, originally prepared for the use of Harrow School, is published in the belief that it may assist the labours of those who are engaged in preparing candidates for Confirmation, and who find it difficult to lay their hand upon any one book of suitable instruction at once sufficiently full to furnish a synopsis of the subject, and sufficiently elastic to give free scope to the individual judgment in the use of it. It will also be found a hand-book for those who are being prepared, as presenting in a compact form the very points which a lecturer would wish his hearers to remember.

2.

The Church Catechism Illustrated and Explained. By
ARTHUR RAMSAY, M.A.
18mo. cloth. 2s.

3.

Hand-Book to Butler's Analogy. By C. A. SWAINSON, M.A. 55 pp. (1856). Crown 8vo. 1s. 6d.

4.

History of the Christian Church during the First Three Centuries, and the Reformation in England. By W. SIMPSON, M.A. Fourth Edition. Fcp. 8vo. cloth. 3s. 6d.

5.

Analysis of Paley's Evidences of Christianity. By CHARLES H. CROSSE, M.A. 115 pp. (1855). 18mo. 3s. 6d.

FORTHCOMING BOOKS.

1.

**An Elementary Treatise on
Natural Philosophy.**

By WILLIAM THOMSON, LL.D., F.R.S., late Fellow of St. Peter's Coll., Cambridge, Professor of Natural Philosophy in the University of Glasgow; and PETER GUTHRIE TAIT, M.A., late Fellow of St. Peter's College, Cambridge, Professor of Natural Philosophy in the University of Edinburgh. With numerous Illustrations.
[In the Press.]

2.

The Narrative of Odysseus.

Homer's Odyssey, Books IX—XII. The Greek Text with English Notes. For Schools and Colleges. By JOHN E. B. MAYOR, M.A., Fellow and Principal Classical Lecturer of St. John's College, Cambridge.
[Nearly Ready.]

3.

**First Book of Algebra. For
Schools.** By J. C. W. ELLIS, M.A., and P. M. CLARKE, M.A., Sidney Sussex College, Cambridge.
[Preparing]

4.

Aristotelis de Rhetorica.

With Notes and Introduction. By E. M. COPE, M.A., Fellow and Assistant Tutor of Trinity College, Cambridge.

5.

The New Testament in the Original Greek. Text revised by B. F. WESTCOTT, M.A., and F. J. HORT, M.A., formerly Fellows of Trinity College.

CAMBRIDGE MANUALS

FOR THEOLOGICAL STUDENTS.

- 1. History of the Christian Church during the Middle Ages.** By ARCHDEACON HARDWICK. Second Edition. 482 pp. (1861). With Maps. Crown 8vo. cloth. 10s. 6d.

This Volume claims to be regarded as an integral and independent treatise on the Mediaeval Church. The History commences with the time of Gregory the Great, to the year 1520,—the year when Luther, having been extruded from those Churches that adhered to the Communion of the Pope, established a provisional form of government and opened a fresh era in the history of Europe.

- 2. History of the Christian Church during the Reformation.** By ARCHD. HARDWICK. 459 pp. (1856). Crown 8vo. cloth. 10s. 6d.

This Work forms a Sequel to the Author's Book on The Middle Ages. The Author's wish has been to give the reader a trustworthy version of those stirring incidents which mark the Reformation period.

- 3. History of the Book of Common Prayer.** With a Rationale of its Offices. By FRANCIS PROCTER, M.A. Fifth Edition. 464 pp. (1860). Crown 8vo. cloth. 10s. 6d.

In the course of the last twenty years the whole question of liturgical knowledge

has been reopened with great learning and accurate research, and it is mainly with the view of epitomizing their extensive publications, and correcting by their help the errors and misconceptions which had obtained currency, that the present volume has been put together.

- 4. History of the Canon of the New Testament during the First Four Centuries.** By BROOKE FOSS WESTCOTT, M.A. 594 pp. (1855). Crown 8vo. cloth. 12s. 6d.

The Author has endeavoured to connect the history of the New Testament Canon with the growth and consolidation of the Church, and to point out the relation existing between the amount of evidence for the authenticity of its component parts and the whole mass of Christian literature. Such a method of inquiry will convey both the truest notion of the connexion of the written Word with the living Body of Christ, and the surest conviction of its divine authority.

- 5. Introduction to the Study of the GOSPELS.** By BROOKE FOSS WESTCOTT, M.A. 458 pp. (1860). Crown 8vo. cloth. 10s. 6d.

This book is intended to be an Introduction to the *Study* of the Gospels. In a subject which involves so vast a literature much must have been overlooked; but the Author has made it a point at least to study the researches of the great writers, and consciously to neglect none.

This Series of THEOLOGICAL MANUALS has been published with the aim of supplying Books concise, comprehensive, and accurate; convenient for the Student, and yet interesting to the general reader.

Uniformly printed $\frac{1}{4}$ 18mo.
with Vignette Titles by
T. Woolner, W. Holman
Hunt, &c.



Handsomely bound in ex-
tra cloth, 4s. 6d. Morocco
plain, 7s. 6d. Morocco ex-
tra, 10s. 6d. Each Volume

1. THE GOLDEN TREASURY OF THE BEST SONGS AND LYRICAL POEMS IN THE ENGLISH LANGUAGE

Selected and arranged, with Notes, by F. T. PALGRAVE.

TWELFTH THOUSAND, with a Vignette by T. WOOLNER.

"There is no book in the English language which will make a more delightful companion than this . . . which must not only be read, but possessed, in order to be adequately valued."—SPECTATOR.

2. THE CHILDREN'S GARLAND. FROM THE BEST POETS.

Selected and Arranged by COVENTRY PATMORE.

FOURTH THOUSAND, with Vignette by T. WOOLNER.

"Mr. Patmore deserves our gratitude for having searched through the wide field of English poetry for these flowers which youth and age can equally enjoy, and woven them into 'The Children's Garland.'"—LONDON REVIEW.

3. THE PILGRIM'S PROGRESS. By JOHN BUNYAN.

With Vignette by W. HOLMAN HUNT.

Large paper copies, crown 8vo. cloth, 7s. 6d., half morocco, 10s. 6d.

"A prettier and better edition and one more exactly suited for use as an elegant and inexpensive Gift Book is not to be found."—EXAMINER.

4. THE BOOK OF PRAISE. FROM THE BEST ENGLISH HYMN WRITERS.

Selected and arranged by ROUNDELL PALMER.

EIGHTH THOUSAND, with Vignette by T. WOOLNER.

"Comprehending nearly all that is excellent in the hymnology of the language. . . . In the details of editorial labours the most exquisite finish is manifest"—THE FREEMAN.

5. BACON'S ESSAYS AND COLOURS OF GOOD AND EVIL.

With Notes and Glossarial Index, by W. ALDIS WRIGHT, M.A.,
Trinity College, Cambridge.

And a Vignette of Woolner's Statue of Lord Bacon.

Large Paper Copies, Crown 8vo. cloth, 7s. 6d., half-morocco, 10s. 6d.

"Edited in a manner worthy of their merit and fame, as an English classic ought to be edited."—DAILY NEWS.

Jonathan Palmer, Printer, Cambridge.

